

# NEET-UG

## Neet 2021 PYQ


2021



## Section -A

1. An inductor of inductance  $L$ , a capacitor of capacitance  $C$  and a resistor of resistance ' $R$ ' are connected in series to an ac source of potential difference ' $V$ ' volts as shown in figure.

Potential difference across  $L$ ,  $C$  and  $R$  is  $40\text{ V}$ ,  $10\text{ V}$  and  $40\text{ V}$ , respectively. The amplitude of current flowing through LCR series circuit is  $I_0 = 10\sqrt{2}\text{ A}$ . The impedance of the circuit is: -




- (A)  $4\sqrt{2}\Omega$   
 (B)  $5/\sqrt{2}\Omega$   
 (C)  $4\Omega$   
 (D)  $5\Omega$

Ans. (D)

Sol.  $I_0 = 10\sqrt{2}\text{ A}$


$$I_{\text{RMS}} = \frac{I_0}{\sqrt{2}} = 10\text{ A}$$



$$\begin{aligned} V_{\text{RMS}} &= \sqrt{V_R^2 + (V_L - V_C)^2} \\ &= \sqrt{(40)^2 + (40 - 10)^2} \\ &= 50\text{ V} \end{aligned}$$

$$Z = \frac{V_{\text{RMS}}}{I_{\text{RMS}}} = \frac{50\text{ V}}{10\text{ A}} = 5\Omega$$


2. Find the value of the angle of emergence from the prism. Refractive index of the glass is  $\sqrt{3}$ .



- (A)  $60^\circ$                       (B)  $30^\circ$                       (C)  $45^\circ$                       (D)  $90^\circ$

Ans. (A)

Sol.




$$r_1 + r_2 = A = 30^\circ \quad \Rightarrow \quad r_2 = 30^\circ \quad (r_1 = 0^\circ)$$

$$\text{from Snell's law} \quad \Rightarrow \quad \sqrt{3} \sin r_2 = 1 \times \sin e$$

$$\sqrt{3} \sin 30^\circ = \sin e$$

$$e = 60^\circ$$

3. A dipole is placed in an electric field as shown. In which direction will it move ?




- (A) towards the left as its potential energy will increase.  
 (B) towards the right as its potential energy will decrease.  
 (C) towards the left as its potential energy will decrease.  
 (D) towards the right as its potential energy will increase.

Ans. (B)

Sol.  $|\vec{E}_1| > |\vec{E}_2|$

as field lines are closer at charge +q, so net force on the dipole acts towards right side. A system always moves to decrease its potential energy.



4. A capacitor of capacitance 'C', is connected across an ac source of voltage V, given by  $V = V_0 \sin \omega t$ . The displacement current between the plates of the capacitor, would then be given by:

(A)  $I_d = V_0 \omega C \cos \omega t$

(B)  $I_d = \frac{V_0}{\omega C} \cos \omega t$

(C)  $I_d = \frac{V_0}{\omega C} \sin \omega t$

(D)  $I_d = V_0 \omega C \sin \omega t$

Ans. (A)





Sol.  $q = CV$

$$\frac{dq}{dt} = \frac{CdV}{dt}$$

$$I_d = C(V_0 \omega \cos \omega t)$$

$$= V_0 \omega C \cos \omega t$$

5. A thick current carrying cable of radius 'R' carries current 'I' uniformly distributed across its cross-section. The variation of magnetic field B(r) due to the cable with the distance 'r' from the axis of the cable is represented by:




Ans. (C)

Sol. Inside a current carrying cylindrical conductor,

$$B = \frac{\mu_0 I}{2\pi R^2} r \quad \therefore B \propto r$$

Outside the conductor,

$$B = \frac{\mu_0 I}{2\pi r} \quad \therefore B \propto \frac{1}{r}$$




6. A convex lens 'A' of focal length 20 cm and a concave lens 'B' of focal length 5 cm are kept along the same axis with a distance 'd' between them. If a parallel beam of light falling on 'A' leaves 'B' as a parallel beam, then the distance 'd' in cm will be :-

(A) 25 (B) 15 (C) 50 (D) 30

Ans. (B)

Sol.



$$\begin{aligned} d &= f_1 - f_2 \\ &= 20 - 5 \\ &= 15 \text{ cm} \end{aligned}$$

7. An electromagnetic wave of wavelength ' $\lambda$ ' is incident on a photosensitive surface of negligible work function. If ' $m$ ' mass is of photoelectron emitted from the surface has de-Broglie wavelength  $\lambda_d$ , then

- (A)  $\lambda = \left(\frac{2m}{hc}\right)\lambda_d^2$   
 (B)  $\lambda_d = \left(\frac{2mc}{h}\right)\lambda^2$   
 (C)  $\lambda = \left(\frac{2mc}{h}\right)\lambda_d^2$   
 (D)  $\lambda = \left(\frac{2h}{mc}\right)\lambda_d^2$

Ans. (C)

Sol.  $\frac{hc}{\lambda} = K_{\max} + \phi$  [given  $\phi$  is negligible]

so,  $\frac{hc}{\lambda} = K_{\max}$

$$\lambda_d = \frac{h}{\sqrt{2mK_{\max}}} \Rightarrow K_{\max} = \frac{h^2}{2m\lambda_d^2}$$

$$\left(\frac{hc}{\lambda}\right) = \frac{h^2}{2m\lambda_d^2} \Rightarrow \lambda = \left(\frac{2mc}{h}\right)\lambda_d^2$$

8. **Column-I** gives certain physical terms associated with flow of current through a metallic conductor. **Column-II** gives some mathematical relations involving electrical quantities. Match **Column-I** and **Column-II** with appropriate relations.

Column-I		Column-II	
(A)	Drift Velocity	(P)	$\frac{m}{ne^2\rho}$
(B)	Electrical Resistivity	(Q)	$nev_d$
(C)	Relaxation Period	(R)	$\frac{eE}{m}\tau$
(D)	Current Density	(S)	$\frac{E}{J}$

(A) (A)-(R), (B)-(S), (C)-(P), (D)-(Q)

(B) (A)-(R), (B)-(S), (C)-(Q), (D)-(P)

(C) (A)-(R), (B)-(P), (C)-(S), (D)-(Q)

(D) (A)-(R), (B)-(Q), (C)-(S), (D)-(P)

Ans. (A)

Sol. (A)  $v_d = \left(\frac{eE}{m}\right)\tau$

(B)  $J = \sigma E = E/\rho$   
 $\Rightarrow \rho = E/J$

(C)  $\rho = \frac{E}{nev_d}$

$v_d = \frac{E}{ne\rho}$

$\frac{eE}{m}\tau = \frac{E}{ne\rho}$

$\tau = \frac{m}{ne^2\rho}$

(D)  $i = neAv_d$

$\frac{i}{A} = nev_d$

$J = nev_d$

9. A radioactive nucleus  ${}^A_ZX$  undergoes spontaneous decay in the sequence  
 ${}^A_ZX \rightarrow {}_{Z-1}B \rightarrow {}_{Z-3}C \rightarrow {}_{Z-2}D$ ,

Where Z is the atomic number of element X. The possible decay particles in the sequence are :

- (A)  $\alpha, \beta^-, \beta^+$       (B)  $\alpha, \beta^+, \beta^-$       (C)  $\beta^+, \alpha, \beta^-$       (D)  $\beta^-, \alpha, \beta^+$

Ans. (C)

Sol.  ${}^A_ZX \xrightarrow{\beta^+} {}_{Z-1}B \xrightarrow{\alpha} {}_{Z-3}C \xrightarrow{\beta^-} {}_{Z-2}D$

$\beta^+$  decreases atomic number by 1

$\alpha$  decreases atomic number by 2

$\beta^-$  increases atomic number by 1

10. The effective resistance of a parallel connection that consists of four wires of equal length, equal area of cross-section and same material is  $0.25 \Omega$ . What will be the effective resistance if they are connected in series ?

(A)  $0.25\Omega$                       (B)  $0.5\Omega$                       (C)  $1\Omega$                       (D)  $4\Omega$


Ans. (D)

Sol.  $R_{\parallel} = \frac{R}{4} = 0.25\Omega$

$$R = 1\Omega$$

$$R_{\text{series}} = 4R$$

$$= 4(1) = 4\Omega$$



11. A particle is released from height  $S$  from the surface of the Earth. At a certain height its kinetic energy is three times its potential energy. The height from the surface of earth and the speed of the particle at that instant are respectively :

(A)  $\frac{S}{4}, \frac{3gS}{2}$

(B)  $\frac{S}{4}, \frac{\sqrt{3gS}}{2}$

(C)  $\frac{S}{2}, \frac{\sqrt{3gS}}{2}$

(D)  $\frac{S}{4}, \sqrt{\frac{3gS}{2}}$

Ans. (D)

Sol.  $U + KE = E$

$$4U = E = mgS$$

$$4mgh = mgS$$

$$h = \frac{S}{4}$$

$$V = \sqrt{2g\left(\frac{3S}{4}\right)} = \sqrt{\frac{3gS}{2}}$$

12. The half-life of a radioactive nuclide is 100 hours. The fraction of original activity that will remain after 150 hours would be :

(A)  $\frac{1}{2}$                       (B)  $\frac{1}{2\sqrt{2}}$                       (C)  $\frac{2}{3}$                       (D)  $\frac{2}{3\sqrt{2}}$

Ans. (B)

Sol.  $\frac{A}{A_0} = \left(\frac{1}{2}\right)^{t/T_{1/2}} = \left(\frac{1}{2}\right)^{150/100} = \frac{1}{2\sqrt{2}}$

13. A cup of coffee cools from  $90^{\circ}\text{C}$  to  $80^{\circ}\text{C}$  in  $t$  minutes, when the room temperature is  $20^{\circ}\text{C}$ . The time taken by a similar cup of coffee to cool from  $80^{\circ}\text{C}$  to  $60^{\circ}\text{C}$  at a room temperature same at  $20^{\circ}\text{C}$  is :

(A)  $\frac{13}{10}t$                       (B)  $\frac{13}{5}t$                       (C)  $\frac{10}{13}t$                       (D)  $\frac{5}{13}t$

Ans. (B)

Sol. According to Newton's law of cooling

$$\frac{T_1 - T_2}{t} = K \left[ \frac{T_1 + T_2}{2} - T_0 \right]$$

For 1<sup>st</sup> cup of coffee,

$$\Rightarrow \frac{90 - 80}{t} = K \left[ \frac{90 + 80}{2} - 20 \right] \dots(1)$$

For 2<sup>nd</sup> cup of coffee,

$$\Rightarrow \frac{80 - 60}{t'} = K \left[ \frac{80 + 60}{2} - 20 \right] \dots(2)$$

Divide (1) by (2)

$$\frac{t'}{2t} = \frac{65}{50} \Rightarrow t' = \frac{13}{5}t$$

14. The number of photons per second on an average emitted by the source of monochromatic light of wavelength 600 nm, when it delivers the power of  $3.3 \times 10^{-3}$  watt will be: ( $h = 6.6 \times 10^{-34}$  Js)

(A)  $10^{18}$                       (B)  $10^{17}$                       (C)  $10^{16}$                       (D)  $10^{15}$

Ans. (C)

Sol.  $p = \frac{nhc}{\lambda} \Rightarrow n = \frac{p\lambda}{hc}$

$$n = \frac{3.3 \times 10^{-3} \times 600 \times 10^{-9}}{6.6 \times 10^{-34} \times 3 \times 10^8} \simeq 10^{16}$$

15. A body is executing simple harmonic motion with frequency 'n', the frequency of its potential energy is:-

(A) n                      (B) 2n                      (C) 3n                      (D) 4n

Ans. (B)

Sol. Displacement equation of SHM of frequency 'n'

$$x = A \sin(\omega t) = A \sin(2\pi n t)$$


Now,

$$\text{Potential energy } U = \frac{1}{2} kx^2 = \frac{1}{2} kA^2 \sin^2(2\pi n t)$$

$$= \frac{1}{2} kA^2 \left[ \frac{1 - \cos(2\pi(2n)t)}{2} \right]$$

So frequency of potential energy = 2n


16. An infinitely long straight conductor carries a current of 5 A as shown. An electron is moving with a speed of  $10^5$  m/s parallel to the conductor. The perpendicular distance between the electron and the conductor is 20 cm at an instant. Calculate the magnitude of the force experienced by the electron at that instant.



- (A)  $4 \times 10^{-20}$  N      (B)  $8\pi \times 10^{-20}$  N      (C)  $4\pi \times 10^{-20}$  N      (D)  $8 \times 10^{-20}$  N

Ans. (D)

Sol.



$$f = ev \left( \frac{\mu_0 i}{2\pi r} \right)$$

$$f = \frac{1.6 \times 10^{-19} \times 10^5 \times 2 \times 10^{-7} \times 5}{0.2}$$

$$f = 8 \times 10^{-20} \text{ Newton}$$

17. If force [F], acceleration [A] and time [T] are chosen as the fundamental physical quantities. Find the dimensions of energy.

- (A) [F] [A] [T]      (B) [F] [A] [T<sup>2</sup>]      (C) [F] [A] [T<sup>-1</sup>]      (D) [F] [A<sup>-1</sup>] [T]

Ans. (B)

Sol.  $E \propto F^a A^b T^c$

$$[M^1 L^2 T^{-2}] \propto [M^1 L^1 T^{-2}]^a [L T^{-2}]^b [T]^c$$

$$a = 1$$

$$a + b = 2 \Rightarrow b = 1$$

$$-2a - 2b + c = -2$$

$$\Rightarrow c = 2$$

$$a = 1 \quad b = 1 \quad c = 2$$

$$E \propto [F][A][T^2]$$

18. Match **Column-I** and **Column-II** and choose the correct match from the given choices.

Column-I		Column-II	
(A)	Root mean square speed of gas molecules	(P)	$\frac{1}{3}nm\bar{v}^2$
(B)	Pressure exerted by ideal gas	(Q)	$\sqrt{\frac{3RT}{M}}$
(C)	Average kinetic energy of a molecule	(R)	$\frac{5}{2}RT$
(D)	Total internal energy of 1 mole of a diatomic gas	(S)	$\frac{3}{2}k_B T$

- (A) (A) - (R), (B) - (P), (C) - (S), (D) - (Q)      (B) (A) - (Q), (B) - (R), (C) - (S), (D) - (P)  
 (C) (A) - (Q), (B) - (P), (C) - (S), (D) - (R)      (D) (A) - (R), (B) - (Q), (C) - (P), (D) - (S)

**Ans.** (C)

**Sol.** Root mean square speed of gas molecules

$$v_{rms} = \sqrt{\frac{3RT}{M}}$$

Pressure exerted by ideal Gas

$$P = \frac{1}{3}\rho v_{rms}^2$$

$$P = \frac{1}{3}nmv^2$$

$$\rho = mn, v_{rms}^2 = \bar{v}^2$$

Average kinetic energy of a molecular

$$KE = \frac{3}{2}KT$$

Total internal energy of 1 mole of a diatomic gas

$$U = \frac{f}{2}\mu RT$$

$$U = \frac{5}{2}RT \quad (\text{For 1 mole diatomic gas})$$

19. A small block slides down on a smooth inclined plane, starting from rest at time  $t = 0$ . Let  $S_n$  be the distance travelled by the block in the interval  $t = n - 1$  to  $t = n$ . Then, the ratio  $\frac{S_n}{S_{n+1}}$  is:

- (A)  $\frac{2n-1}{2n}$       (B)  $\frac{2n-1}{2n+1}$       (C)  $\frac{2n+1}{2n-1}$       (D)  $\frac{2n}{2n-1}$

**Ans.** (B)

**Sol.**  $S_n$  = Distance in  $n^{\text{th}}$  sec. i.e.  $t = n - 1$  to  $t = n$

$S_{n+1}$  = Distance in  $(n + 1)^{\text{th}}$  sec.

i.e.  $t = n$  to  $t = n + 1$

So as we know


$$S_n = \frac{a}{2}(2n-1) \quad a = \text{acceleration}$$

$$\frac{S_n}{S_{n+1}} = \frac{\frac{a}{2}(2n-1)}{\frac{a}{2}(2(n+1)-1)} = \frac{2n-1}{2n+1}$$

$$\frac{S_n}{S_{n+1}} = \frac{2n-1}{2n+1}$$

20. A nucleus with mass number 240 breaks into two fragments each of mass number 120, the binding energy per nucleon of unfragmented nuclei is 7.6 MeV while that of fragments is 8.5 MeV. The total gain in the Binding Energy in the process is :
- (A) 0.9 MeV                      (B) 9.4 MeV                      (C) 804 MeV                      (D) 216 MeV

Ans. (D)



given binding energy per nucleon of X, Y & Z are 7.6 MeV, 8.5 MeV & 8.5 MeV respectively. Gain in binding energy is :-

$$\begin{aligned} Q &= \text{Binding Energy of products} - \text{Binding energy of reactants} \\ &= (120 \times 8.5 \times 2) - (240 \times 7.6) \text{ MeV} \\ &= 216 \text{ MeV} \end{aligned}$$

21. A screw gauge gives the following readings when used to measure the diameter of a wire Main scale reading: 0 mm  
Circular scale reading: 52 divisions  
Given that 1 mm on main scale corresponds to 100 divisions on the circular scale. The diameter of the wire from the above data is:
- (A) 0.52 cm                      (B) 0.026 cm                      (C) 0.26 cm                      (D) 0.052 cm


Ans. (D)

Sol. Least count =  $\frac{1\text{mm}}{100} = 0.01\text{mm}$

Diameter = main scale reading + least count  $\times$  circular scale reading

$$\begin{aligned} \text{Diameter} &= 0 + 52 \times 0.01 \text{ mm} \\ &= 0.52 \text{ mm} = 0.052 \text{ cm} \end{aligned}$$


22. The equivalent capacitance of the combination shown in the figure is :



- (A) 3C                      (B) 2C                      (C) C/2                      (D) 3C/2

Ans. (B)

Sol.



(one capacitor gets short)

$$\begin{aligned} \Rightarrow C_{\text{eq}} &= C_1 + C_2 \\ &= C + C \\ &= 2C \end{aligned}$$

23. A lens of large focal length and large aperture is best suited as an objective of an astronomical telescope since :
- (A) a large aperture contributes to the quality and visibility of the images.  
 (B) a large area of the objective ensures better light gathering power.  
 (C) a large aperture provides a better resolution.  
 (D) all of the above.

Ans. (D)

Sol.  $MP = \frac{f_0}{f_e}$

$$R.P. = \frac{a}{1.22\lambda}$$

large aperture(a) of the objective lens provides better resolution

$\therefore$  good quality of image is formed and also it gathers more light.

24. Two charged spherical conductors of radius  $R_1$  and  $R_2$  are connected by a wire. Then the ratio of surface charge densities of the spheres ( $\sigma_1/\sigma_2$ ) is:

(A)  $\frac{R_1}{R_2}$

(B)  $\frac{R_2}{R_1}$

(C)  $\sqrt{\left(\frac{R_1}{R_2}\right)}$

(D)  $\frac{R_1^2}{R_2^2}$

Ans. (B)

Sol. For a conducting sphere

$$E = \frac{\sigma}{\epsilon_0}$$

$$V = \frac{\sigma R}{\epsilon_0}$$

as both spheres have same potential after connecting with wire,

$$V_1 = V_2$$

$$\sigma_1 R_1 = \sigma_2 R_2$$

$$\Rightarrow \frac{\sigma_1}{\sigma_2} = \frac{R_2}{R_1}$$

25. A spring is stretched T by 5 cm by a force 10 N. The time period of the oscillations when a mass of 2 kg is suspended by it is :

(A) 0.0628 s

(B) 6.28 s

(C) 3.14 s

(D) 0.628 s

Ans. (D)

Sol.  $F = kx$

$$10 = k(5 \times 10^{-2})$$

$$k = \frac{10}{5 \times 10^{-2}} = 2 \times 10^2 = 200 \text{ N/m}$$

$$\text{Now } T = 2\pi \sqrt{\frac{m}{k}} = 2\pi \sqrt{\frac{2}{200}} = \frac{2\pi}{10} = 0.628 \text{ sec.}$$

26. For a plane electromagnetic wave propagating in x-direction, which one of the following combinations gives the correct possible directions for electric field (E) and magnetic field (B) respectively?

- (A)  $\hat{j} + \hat{k}, \hat{j} + \hat{k}$  (B)  $-\hat{j} + \hat{k}, -\hat{j} - \hat{k}$   
 (C)  $\hat{j} + \hat{k}, -\hat{j} - \hat{k}$  (D)  $-\hat{j} + \hat{k}, -\hat{j} + \hat{k}$

Ans. (B)

Sol.  $\vec{v} \parallel \vec{E} \times \vec{B}, \hat{v} = \hat{i}$

Option (A)  $\vec{E} \times \vec{B} = \vec{0} (\vec{E} \parallel \vec{B})$

Option (B)  $\vec{E} \times \vec{B} = 2\hat{i}$  (parallel to  $\vec{v}$ )

Option (C)  $\vec{E} \times \vec{B} = \vec{0} (\vec{E} \uparrow \downarrow \vec{B})$

Option (D)  $\vec{E} \times \vec{B} = \vec{0} (\vec{E} \parallel \vec{B})$

27. The escape velocity from the Earth's surface is  $v$ . The escape velocity from the surface of another planet having a radius, four times that of Earth and same mass density is:

- (A)  $v$  (B)  $2v$  (C)  $3v$  (D)  $4v$

Ans. (D)

Sol.  $v_e = \sqrt{\frac{2GM}{R}} = \sqrt{\frac{2G}{R} \times \frac{4}{3} \pi R^3 \rho}$   
 $= \sqrt{\frac{8\pi G \rho}{3} R^2}$   
 $\Rightarrow v_e \propto R$   
 $\Rightarrow \frac{v_e}{v} = \frac{4R}{R} \Rightarrow v_e = 4v$

28. In a potentiometer circuit a cell of EMF 1.5 V gives balance point at 36 cm length of wire. If another cell of EMF 2.5 V replaces the first cell, then at what length of the wire, the balance point occurs?

- (A) 60 cm (B) 21.6 cm (C) 64 cm (D) 62 cm

Ans. (A)

Sol.  $\frac{E_1}{E_2} = \frac{\phi \ell_1}{\phi \ell_2}$


$$\frac{1.5}{2.5} = \frac{36}{\ell_2} \Rightarrow \ell_2 = 36 \times \frac{5}{3} = 60 \text{ cm}$$

30. The velocity of a small ball of mass  $M$  and density  $d$ , when dropped in a container filled with glycerine becomes constant after some time. If the density of glycerine is  $\frac{d}{2}$ , then the viscous force acting on the ball will be:

(A)  $\frac{Mg}{2}$                       (B)  $Mg$                       (C)  $\frac{3}{2}Mg$                       (D)  $2Mg$

Ans. (A)

Sol. Mass =  $M$



Density of ball =  $d$

Density of glycerine =  $\frac{d}{2}$

$$F_B = V_s \rho_l g = V \frac{d}{2} g$$

$$F_g = mg = vdg$$

for constant velocity,  $F_{\text{net}} = 0$

$$\therefore F_B + F_v = M_g$$

$$F_v = M_g - F_B = Vdg - \frac{Vdg}{2} = \frac{Vdg}{2} = \frac{Mg}{2}$$

30. A parallel plate capacitor has a uniform electric field ' $\vec{E}$ ' in the space between the plates. If the distance between the plates is ' $d$ ' and the area of each plate is ' $A$ ', the energy stored in the capacitor is : ( $\epsilon_0$  = permittivity of free space)

(A)  $\frac{1}{2}\epsilon_0 E^2$                       (B)  $\epsilon_0 EAd$                       (C)  $\frac{1}{2}\epsilon_0 E^2 Ad$                       (D)  $\frac{E^2 Ad}{\epsilon_0}$

Ans. (C)

Sol.  $E = \frac{1}{2} CV^2$

$$= \frac{1}{2} \left( \frac{\epsilon_0 A}{d} \right) (Ed)^2$$

$$= \frac{1}{2} \epsilon_0 E^2 Ad$$

31. The electron concentration in an n-type semiconductor is the same as hole concentration in a p-type semiconductor. An external field (electric) is applied across each of them. Compare the currents in them.
- (A) current in n-type = current in p-type  
 (B) current in p-type > current in n-type  
 (C) current in n-type > current in p-type  
 (D) No current will flow in p-type, current will only flow in n-type

Ans. (C)

Sol. In N type semiconductor majority charge carriers are  $e^-$  and P type semiconductor majority charge carriers are holes.

$$I = neAV_d = neA(\mu E)$$

$$\mu_e > \mu_h \Rightarrow I_e > I_h$$

32. Consider the following statements (A) and (B) and identify the correct answer.
- (A) A zener diode is connected in reverse bias, when used as a voltage regulator.  
 (B) The potential barrier of p-n junction lies between 0.1 V to 0.3 V.
- (A) (A) and (B) both are correct. (B) (A) and (B) both are incorrect.  
 (C) (A) is correct and (B) is incorrect. (D) (A) is incorrect but (B) is correct.

Ans. (C)

Sol. Reverse bias Zener diode use as a voltage regulator

for Ge Potential barrier  $V_0 = 0.3$  V

Si Potential barrier  $V_0 = 0.7$  V

33. Polar molecules are the molecules :
- (A) having zero dipole moment.  
 (B) acquire a dipole moment only in the presence of electric field due to displacement of charges.  
 (C) acquire a dipole moment only when magnetic field is absent.  
 (D) having a permanent electric dipole moment.

Ans. (D)

Sol. Polar molecules have centres of positive and negative charges separated by some distance, so they have permanent dipole moment.

34. If E and G respectively denote energy and gravitational constant, then  $\frac{E}{G}$  has the dimensions of
- (A)  $[M^2][L^{-1}][T^0]$  (B)  $[M][L^{-1}][T^{-1}]$  (C)  $[M][L^0][T^0]$  (D)  $[M^2][L^{-2}][T^{-1}]$

Ans. (A)

Sol.  $E = \text{energy} = [ML^2T^{-2}]$

$G = \text{Gravitational constant} = [M^{-1}L^3T^{-2}]$

$$\text{So } \frac{E}{G} = \frac{[E]}{[G]} = \frac{ML^2T^{-2}}{M^{-1}L^3T^{-2}} = [M^2L^{-1}T^0]$$

Q.35 Water falls from a height of 60 m at the rate of 15 kg/s to operate a turbine. The losses due to frictional force are 10% of the input energy. How much power is generated by the turbine? ( $g = 10 \text{ m/s}^2$ )

- (A) 10.2 kW                      (B) 8.1 kW                      (C) 12.3 kW                      (D) 7.0 kW

Sol. (B)

$$P_{\text{in}} = \frac{mgh}{t} = \frac{15 \times 10 \times 60}{1} = 9000 \text{ w}$$

$$P_{\text{out}} = 90\% \text{ of } P_{\text{in}}$$

$$\Rightarrow 8.1 \text{ kw}$$

### SECTION-B

Q.36 A car starts from rest and accelerates at  $5 \text{ m/s}^2$ . At  $t = 4 \text{ s}$ , a ball is dropped out of a window by a person sitting in the car. What is the velocity and acceleration of the ball at  $t = 6 \text{ s}$ ? (Take  $g = 10 \text{ m/s}^2$ )

- (A) 20 m/s,  $5 \text{ m/s}^2$                       (B) 20 m/s, 0  
(C)  $20\sqrt{2} \text{ m/s}$ , 0                      (D)  $20\sqrt{2} \text{ m/s}$ ,  $10 \text{ m/s}^2$

Sol. (4)

Velocity of car at  $t = 4 \text{ sec}$  is

$$v = u + at$$

$$v = 0 + 5(4)$$

$$= 20 \text{ m/s}$$

At  $t = 6 \text{ sec}$

acceleration is due to gravity  $\square \square a = g = 10 \text{ m/s}^2$

$v_x = 20 \text{ m/s}$  (due to car)

$$v_y = u + at$$

$$= 0 + g(2) \text{ (downward)}$$





$$= 20 \text{ m/s (downward)}$$

$$v = \sqrt{v_x^2 + v_y^2}$$

$$= \sqrt{20^2 + 20^2}$$

$$= 20\sqrt{2} \text{ m/s}$$

Q.37 For the given circuit, the input digital signals are applied at the terminals A, B and C. What would be the output at the terminal y ?



Sol.

(C)

$$Y = A \cdot B + \overline{B} \cdot \overline{C}$$

(i) 0 to  $t_1$      $A = 0, B = 0, C = 1$

$$Y = 0 \cdot 0 + \overline{0} \cdot \overline{1} = 0 + 1 = 1$$

(ii)  $t_1$  to  $t_2$      $A = 1, B = 0, C = 1$

$$Y = 1 \cdot 0 + \overline{0} \cdot \overline{1} = 0 + 1 = 1$$

(iii)  $t_2$  to  $t_3$      $A = 0, B = 1, C = 0$

$$Y = 0 \cdot 1 + \overline{1} \cdot \overline{0} = 0 + 1 = 1$$

Q.38 A ball of mass 0.15 kg is dropped from a height 10 m, strikes the ground and rebounds to the same height. The magnitude of impulse imparted to the ball is ( $g = 10 \text{ m/s}^2$ ) nearly :

- (A) 0 kg m/s      (B) 4.2 kg m/s      (C) 2.1 kg m/s      (D) 1.4 kg m/s

Sol. (B)

Velocity just before striking the ground

$$v_1 = \sqrt{2gh}$$

$$v_1 = \sqrt{2(10)(10)} = 10\sqrt{2} \text{ m/s}$$

$$v_1 = -10\sqrt{2}\hat{j}$$

If it reaches the same height, speed remains same after collision only the direction changes.

$$v_2 = 10\sqrt{2} \text{ m/s}$$

$$\vec{v}_2 = 10\sqrt{2}\hat{j}$$

$$|\text{Impulse}| = m |\Delta \vec{v}|$$

$$= m |10\sqrt{2}\hat{j} - (-10\sqrt{2}\hat{j})|$$


$$= 0.15 [2(10\sqrt{2})]$$

$$= 3\sqrt{2} \text{ kg m/s}$$

$$= 4.2 \text{ kg m/s}$$

Q.39 A uniform rod of length 200 cm and mass 500 g is balanced on a wedge placed at 40 cm mark. A mass of 2 kg is suspended from the rod at 20 cm and another unknown mass 'm' is suspended from the rod at 160 cm mark as shown in the figure. Find the value of 'm' such that the rod is in equilibrium.

( $g = 10 \text{ m/s}^2$ )




(A)  $\frac{1}{2}$  kg

(B)  $\frac{1}{3}$  kg

(C)  $\frac{1}{6}$  kg

(D)  $\frac{1}{12}$  kg

Sol. (4)




By balancing torque

$$2g \times 20 = 0.5g \times 60 + mg \times 120$$

$$m = \frac{0.5}{6} \text{ kg} = \frac{1}{12} \text{ kg}$$

- Q.40 A point object is placed at a distance of 60 cm from a convex lens of focal length 30 cm. If a plane mirror were put perpendicular to the principal axis of the lens and at a distance of 40 cm from it, the final image would be formed at a distance of :



- (A) 20 cm from the lens, it would be a real image.  
 (B) 30 cm from the lens, it would be a real image.  
 (C) 30 cm from the plane mirror, it would be a virtual image.  
 (D) 20 cm from the plane mirror, it would be a virtual image.

Sol.


(4)  
 first, for image formation from lens

$$u = -60 \text{ cm}$$

$$f = +30 \text{ cm}$$

$$\Rightarrow v = \frac{uf}{u+f} = \frac{-60 \times 30}{-60+30} = 60 \text{ cm}$$

this real image formed by lens acts as virtual object for mirror




Real image from plane mirror is formed 20 cm in front of mirror, hence at 20 cm distance from lens. Now, for second refraction from lens,

$$u = -20 \text{ cm}$$

$$f = +30 \text{ cm}$$

$$v = \frac{uf}{u+f} = \frac{-20 \times 30}{-20+30} = -60 \text{ cm}$$

So, final virtual image is 60 cm from lens, or 20 cm behind mirror




- Q.41 A step down transformer connected to an ac mains supply of 220 V is made to operate at 11V, 44 W lamp. Ignoring power losses in the transformer, what is the current in the primary circuit ?  
 (A) 0.2 A (B) 0.4 A (C) 2A (D) 4A

Sol. (1)

$$220 \times i_p = 44$$

$$\Rightarrow i_p = \frac{44}{220} = \frac{1}{5} = 0.2A$$

- Q.42 Three resistors having resistances  $r_1$ ,  $r_2$  and  $r_3$  are connected as shown in the given circuit. The ratio  $\frac{i_3}{i_1}$  of currents in terms of resistances used in the circuit is:



(A)  $\frac{r_1}{r_2 + r_3}$

(B)  $\frac{r_2}{r_2 + r_3}$

(C)  $\frac{r_1}{r_1 + r_2}$

(D)  $\frac{r_2}{r_1 + r_3}$

Sol. (2)

$$V = ir$$


$$i = \frac{V}{r}$$

$$i \propto \frac{1}{r} \quad [v \text{ is same for } r_2 \& r_3]$$

$$\frac{i_2}{i_3} = \frac{r_3}{r_2}$$

$$i_3 = \frac{r_2}{r_2 + r_3} i_1$$

$$\frac{i_3}{i_1} = \frac{r_2}{r_2 + r_3}$$



Q.43 In the product

$$\vec{F} = q(\vec{v} \times \vec{B})$$

$$= q\vec{v} \times (B_1\hat{i} + B_2\hat{j} + B_0\hat{k})$$

For  $q = 1$  and  $\vec{v} = 2\hat{i} + 4\hat{j} + 6\hat{k}$  and  $\vec{F} = 4\hat{i} - 20\hat{j} + 12\hat{k}$

What will be the complete expression for  $\vec{B}$ ?

- (A)  $-8\hat{i} - 8\hat{j} - 6\hat{k}$       (B)  $-6\hat{i} - 6\hat{j} - 8\hat{k}$       (C)  $8\hat{i} + 8\hat{j} - 6\hat{k}$       (D)  $6\hat{i} + 6\hat{j} - 8\hat{k}$

Sol. (2)

$$\vec{F} = q(\vec{v} \times \vec{B})$$

$$4\hat{i} - 20\hat{j} + 12\hat{k} = 1 \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 4 & 6 \\ B & B & B_0 \end{vmatrix}$$

Comparing

$$\begin{cases} 4 = 4B_0 - 6B \\ \Rightarrow -20 = -2B_0 + 6B \\ 12 = 2B - 4B \end{cases} \begin{matrix} \text{Solving} \\ B = -6 \\ B_0 = -8 \end{matrix}$$

$$\therefore \vec{B} = -6\hat{i} - 6\hat{j} - 8\hat{k}$$

Q.44 A particle of mass 'm' is projected with a velocity  $v = kV_e$  ( $k < 1$ ) from the surface of the earth. ( $V_e$  = escape velocity) The maximum height above the surface reached by the particle is :

- (A)  $R \left( \frac{k}{1-k} \right)^2$       (B)  $R \left( \frac{k}{1+k} \right)^2$       (C)  $\frac{R^2 k}{1+k}$       (D)  $\frac{Rk^2}{1-k^2}$

Sol. (4)

$$h = \frac{R}{\frac{2gR}{V^2} - 1} = \frac{R}{\frac{V_e^2}{K^2 V_e^2} - 1} = \frac{RK^2}{1-K^2}$$

Q.45 Twenty seven drops of same size are charged at 220 V each. They combine to form a bigger drop. Calculate the potential of the bigger drop.

- (A) 660 V      (B) 1320 V      (C) 1520 V      (D) 1980 V

Sol. (D)

$$\frac{4}{3}\pi R^3 = 27 \left( \frac{4}{3}\pi r^3 \right) \Rightarrow R = 3r \quad \dots (1)$$

$$V = \frac{Kq}{r} \Rightarrow \frac{V_1}{V_2} = \left( \frac{q_1}{q_2} \right) \left( \frac{r_2}{r_1} \right)$$

$$\Rightarrow \frac{220}{V_2} = \left( \frac{q}{27q} \right) \left( \frac{3r}{r} \right)$$

$$\Rightarrow \frac{220}{V_2} = \frac{1}{9}$$

$$\Rightarrow V_2 = 220 \times 9 = 1980 \text{ Volt}$$

- Q.46 A series LCR circuit containing 5.0 H inductor, 80  $\mu\text{F}$  capacitor and 40  $\Omega$  resistor is connected to 230 V variable frequency ac source. The angular frequencies of the source at which power transferred to the circuit is half the power at the resonant angular frequency are likely to be :
- (A) 25 rad/s and 75 rad/s (B) 50 rad/s and 25 rad/s  
(C) 46 rad/s and 54 rad/s (D) 42 rad/s and 58 rad/s

Sol. (C)

$$Q = \frac{\omega}{\Delta\omega} = \frac{\omega L}{R} \Rightarrow \Delta\omega = R/L = \frac{50}{4} = 8 \text{ rad/sec}$$

$$\omega_0 = \frac{1}{\sqrt{LC}} = \frac{1}{\sqrt{5 \times 80 \times 10^{-6}}} = 50 \text{ rad/sec.}$$

$$\omega_{\min} = \omega_0 - \frac{\Delta\omega}{2} = 46 \text{ rad/sec}$$

$$\omega_{\max} = \omega_0 + \frac{\Delta\omega}{2} = 54 \text{ rad/sec}$$

- Q.47 A uniform conducting wire of length 12a and resistance 'R' is wound up as a current carrying coil in the shape of,
- (i) an equilateral triangle of side 'a'.  
(ii) a square of side 'a'.

The magnetic dipole moments of the coil in each case respectively are:

- (A)  $\sqrt{3} I a^2$  and  $3 I a^2$  (B)  $3 I a^2$  and  $I a^2$   
(C)  $3 I a^2$  and  $4 I a^2$  (D)  $4 I a^2$  and  $3 I a^2$

Sol. (A)

$$M_1 = \left( \frac{\sqrt{3}}{4} a^2 \right) I \times 4 = \sqrt{3} I a^2 \text{ (no. of turns = 4)}$$

$$M_2 = a^2 I \times 3 = 3 I a^2 \text{ (no. of turns = 3)}$$

- Q.48 From a circular ring of mass 'M' and radius 'R' an arc corresponding to a 90° sector is removed. The moment of inertia of the remaining part of the ring about an axis passing through the centre of the ring and perpendicular to the plane of the ring is 'K' times 'MR<sup>2</sup>'. Then the value of 'K' is :


- (A)  $\frac{3}{4}$  (B)  $\frac{7}{8}$  (C)  $\frac{1}{4}$  (D)  $\frac{1}{8}$

Sol. (A)

$$M_{\text{remain}} = \frac{3}{4} M$$

$$I = M_{\text{remain}} R^2$$

$$= \frac{3}{4} M R^2$$



Q.49 Two conducting circular loops of radii  $R_1$  and  $R_2$  are placed in the same plane with their centres coinciding. If  $R_1 \gg R_2$ , the mutual inductance  $M$  between them will be directly proportional to:


- (A)  $\frac{R_1}{R_2}$                       (B)  $\frac{R_2}{R_1}$                       (C)  $\frac{R_1^2}{R_2}$                       (D)  $\frac{R_2^2}{R_1}$

Sol. (D)

$$M = \frac{\phi_{12}}{I_1} = \frac{B_1 A_2}{I_1} = \frac{\left(\frac{\mu_0 I_1}{2R_1}\right) (\pi R_2^2)}{I_1}$$

$$M = \frac{\mu_0 \pi R_2^2}{2R_1}$$

$$M \propto \frac{R_2^2}{R_1}$$



Q.50 A particle moving in a circle of radius  $R$  with a uniform speed takes a time  $T$  to complete one revolution.

If this particle were projected with the same speed at an angle ' $\theta$ ' to the horizontal, the maximum height attained by it equals  $4R$ . The angle of projection,  $\theta$ , is then given by :

(A)  $\theta = \cos^{-1} \left( \frac{gT^2}{\pi^2 R} \right)^{1/2}$

(B)  $\theta = \cos^{-1} \left( \frac{\pi^2 R}{gT^2} \right)^{1/2}$

(C)  $\theta = \sin^{-1} \left( \frac{\pi^2 R}{gT^2} \right)^{1/2}$

(D)  $\theta = \sin^{-1} \left( \frac{2gT^2}{\pi^2 R} \right)^{1/2}$

Sol. (D)

$$T = \frac{2\pi R}{v} \Rightarrow v = \frac{2\pi R}{T} \quad \dots(1)$$

$$H_{\max} = \frac{v^2 \sin^2 \theta}{2g} = \frac{2\pi^2 R^2 \sin^2 \theta}{gT^2} = 4R$$

$$\sin \theta = \left( \frac{2gT^2}{\pi^2 R} \right)^{1/2}$$

$$\theta = \sin^{-1} \left[ \frac{2gT^2}{\pi^2 R} \right]^{1/2}$$

## SECTION-B

51. Given below are two statements:

**Statement I:** Aspirin and Paracetamol belong to the class of narcotic analgesics.




**Statement II:** Morphine and Heroin are non-narcotic analgesics. In the light of the above statements, choose the correct answer from the options given below.

- (A) Both Statement I and Statement II are true  
 (B) Both Statement I and Statement II are false  
 (C) Statement I is correct but Statement II is false  
 (D) Statement I is incorrect but Statement II is true.


Ans. (B)

Sol. • Aspirin and paracetamol belong to the class of non-narcotic analgesics  
 • Morphine and Heroin are Narcotic analgesics  
 ∴ Both statement I and statement II are false

52. The correct structure of 2, 6-Dimethyl-dec-4-ene is



Ans. (A)




2, 6-Dimethyldec-4-ene

53.  $\text{BF}_3$  is planar and electron deficient compound. Hybridization and number of electrons around the central atom, respectively are :

- (A)  $\text{sp}^3$  and 4                      (B)  $\text{sp}^3$  and 6                      (C)  $\text{sp}^2$  and 6                      (D)  $\text{sp}^2$  and 8

Ans. (C)



- Number of electrons around boron atom is 6.
- Hybridization of B is  $\text{sp}^2$ .
- Shape is trigonal planar.

54. Noble gases are named because of their inertness towards reactivity. Identify an incorrect statement about them.

- (A) Noble gases are sparingly soluble in water  
 (B) Noble gases have very high melting and boiling points  
 (C) Noble gases have weak dispersion forces  
 (D) Noble gases have large positive values of electron gain enthalpy

Ans. (B)

Sol. Noble gases have weak dispersion forces hence they have low melting and boiling points.

55. The molar conductance of  $\text{NaCl}$ ,  $\text{HCl}$  and  $\text{CH}_3\text{COONa}$  at infinite dilution are 126.45, 426.16 and  $91.0 \text{ Scm}^2\text{mol}^{-1}$  respectively. The molar conductance of  $\text{CH}_3\text{COOH}$  at infinite dilution is. Choose the right option for your answer.

- (A)  $201.28 \text{ Scm}^2\text{mol}^{-1}$   
 (B)  $390.71 \text{ Scm}^2\text{mol}^{-1}$   
 (C)  $698.28 \text{ Scm}^2\text{mol}^{-1}$   
 (D)  $540.48 \text{ Scm}^2\text{mol}^{-1}$

Ans. (B)

Sol. According to Kohlrausch law of independent migration of ions.

$$\begin{aligned} \Lambda_m^\circ(\text{CH}_3\text{COOH}) &= \Lambda_m^\circ(\text{CH}_3\text{COONa}) + \Lambda_m^\circ(\text{HCl}) - \Lambda_m^\circ(\text{NaCl}) \\ &= 91.0 \text{ Scm}^2\text{mol}^{-1} + 426.16 \text{ Scm}^2\text{mol}^{-1} - 126.45 \text{ Scm}^2\text{mol}^{-1} \\ &= 390.71 \text{ Scm}^2\text{mol}^{-1} \end{aligned}$$

56. The right option for the statement "Tyndall effect is exhibited by", is :

- (A)  $\text{NaCl}$  solution  
 (B) Glucose solution  
 (C) Starch solution  
 (D) Urea solution

Ans. (C)

Sol. • Tyndall effect is exhibited by colloidal solution only.

- Among the given options, Urea,  $\text{NaCl}$  and Glucose solutions are true solutions, so cannot show Tyndall effect.
- Starch solution is a colloidal solution therefore can show Tyndall effect.

57. The RBC deficiency is deficiency disease of :

- (A) Vitamin B<sub>12</sub>
- (B) Vitamin B<sub>6</sub>
- (C) Vitamin B<sub>1</sub>
- (D) Vitamin B<sub>2</sub>

Ans. (A)

- Sol.
- Deficiency of vitamin B<sub>2</sub> (Riboflavin) causes cheilosis, digestive disorders and burning sensation of the skin.
  - Deficiency of vitamin B<sub>12</sub> causes Pernicious anaemia which is RBC deficiency in haemoglobin.
  - Deficiency of vitamin B<sub>6</sub> (Pyridoxine) causes Convulsions.
  - Deficiency of vitamin B<sub>1</sub> (Thiamine) causes Beri-Beri (loss of appetite and retarded growth).

58. Dihedral angle of least stable conformer of ethane is :


- (A) 120°
- (B) 180°
- (C) 60°
- (D) 0°

Ans. (D)

Sol. Ethane has two conformers

- (i) Eclipsed
- (ii) Staggered

Eclipsed conformer is least stable while staggered conformer is most stable. In eclipsed conformer the dihedral angle is 0°



59. The incorrect statement among the following is :

- (A) Actinide contraction is greater for element to element than lanthanide contraction
- (B) Most of the trivalent lanthanide ions are colorless in the solid state
- (C) Lanthanides are good conductors of heat and electricity
- (D) Actinides are highly reactive metals, especially when finely divided.


Ans. (B)

- Sol.
- Actinides are highly reactive metals, especially when finely divided
  - Actinide contraction is greater from element to element than lanthanide contraction resulting from poor shielding by 5f electrons
  - Many trivalent lanthanides ions are coloured both in the solid state and in aqueous solutions.
  - Lanthanides have typical metallic structure and are good conductors of heat and electricity

60. The major product formed in dehydrohalogenation reaction of 2-Bromo pentane is Pent-2-ene. This product formation is based on?  
 (A) Saytzeff's Rule (B) Hund's Rule  
 (C) Hofmann Rule (D) Huckel's Rule

**Ans. (A)**

**Sol.** Major product formed in dehydrohalogenation reaction of 2-bromopentane is pent-2-ene because according to Saytzeff's rule, in dehydrohalogenation reactions, the preferred product is that alkene which has greater number of alkyl group(s) attached to the doubly bonded carbon atoms.



61. Which one among the following is the correct option for right relationship between  $C_P$  and  $C_V$  for one mole of ideal gas?  
 (1)  $C_P + C_V = R$   
 (B)  $C_P - C_V = R$   
 (C)  $C_P = RC_V$   
 (D)  $C_V = RC_P$


**Ans. (B)**

**Sol.** At constant volume,  $q_V = C_V \Delta T = \Delta U$   
 At constant pressure,  $q_P = C_P \Delta T = \Delta H$   
 For a mole of an ideal gas,  
 $\Delta H = \Delta U + \Delta(PV)$   
 $= \Delta U + \Delta(RT)$   
 $= \Delta U + R\Delta T$   
 On putting the values of  $\Delta H$  and  $\Delta U$ , we have  
 $C_P \Delta T = C_V \Delta T + R\Delta T$   
 $C_P = C_V + R$   
 $C_P - C_V = R$


62. Which one of the following polymers is prepared by addition polymerisation?  
 (A) Teflon (B) Nylon-6, 6 (C) Novolac (D) Dacron

**Ans. (A)**

**Sol.** Dacron, Nylon-66 and Novolac are prepared by condensation polymerisation. Teflon is an addition polymer. Monomer of teflon is tetrafluoroethene.



63. What is the IUPAC name of the organic compound formed in the following chemical reaction?




(A) 2-methylpropan-2-ol

(B) pentan-2-ol

(C) pentan-3-ol

(D) 2-methylbutan-2-ol

Ans. (D)



64. Match List-I with List-II.

**List-I**

(a)  $\text{PCl}_5$

(b)  $\text{SF}_6$

(c)  $\text{BrF}_5$

(d)  $\text{BF}_3$

**List-II**

(i) Square pyramidal

(ii) Trigonal planar

(iii) Octahedral

(iv) Trigonal bipyramidal

Choose the correct answer from the options given below.


(A) (a)-(iv), (b)-(iii), (c)-(i), (d)-(ii)

(B) (a)-(ii), (b)-(iii), (c)-(iv), (d)-(i)

(C) (a)-(iii), (b)-(i), (c)-(iv), (d)-(ii)

(D) (a)-(iv), (b)-(iii), (c)-(ii), (d)-(i)

Ans. (A)




65. Which one of the following methods can be used to obtain highly pure metal which is liquid at room temperature?

- (A) Electrolysis      (B) Chromatography      (C) Distillation      (D) Zone refining

Ans. (C)


Sol. Distillation method is generally used for the purification of metals having low boiling point such as Hg, Zn etc.

66. The major product of the following chemical reaction is :







- (A)  $\begin{array}{c} \text{CH}_3 \\ | \\ \text{CH}-\text{CH}_2-\text{CH}_2-\text{Br} \\ | \\ \text{CH}_3 \end{array}$       (B)  $\begin{array}{c} \text{CH}_3 \\ | \\ \text{CH}-\text{CH}_2-\text{CH}_2-\text{O}-\text{COC}_6\text{H}_5 \\ | \\ \text{CH}_3 \end{array}$
- (C)  $\begin{array}{c} \text{CH}_3 \\ | \\ \text{CH}-\text{CH}-\text{CH}_3 \\ | \\ \text{CH}_3 \\ | \\ \text{Br} \end{array}$       (D)  $\begin{array}{c} \text{CH}_3 \\ | \\ \text{CBr}-\text{CH}_2-\text{CH}_3 \\ | \\ \text{CH}_3 \end{array}$

Ans. (A)



**Mechanism :** Peroxide effect proceeds via free radical chain mechanism.



67. Tritium, a radioactive isotope of hydrogen, emits which of the following particles?  
 (A) Beta ( $\beta^-$ )                      (B) Alpha ( $\alpha$ )                      (C) Gamma ( $\gamma$ )                      (D) Neutron (n)

Ans. (A)

Sol. Hydrogen has three isotopes : protium or  ${}^1_1\text{H}$ , deuterium  ${}^2_1\text{H}$  or D and tritium  ${}^3_1\text{H}$  or T. Of these isotopes, only tritium is radioactive and emits low energy  $\beta^-$  particles ( $t_{1/2} = 12.33$  years).

68. The correct sequence of bond enthalpy of 'C-X' bond is :

- (A)  $\text{CH}_3 - \text{F} < \text{CH}_3 - \text{Cl} < \text{CH}_3 - \text{Br} < \text{CH}_3 - \text{I}$   
 (B)  $\text{CH}_3 - \text{F} > \text{CH}_3 - \text{Cl} > \text{CH}_3 - \text{Br} > \text{CH}_3 - \text{I}$   
 (C)  $\text{CH}_3 - \text{F} < \text{CH}_3 - \text{Cl} > \text{CH}_3 - \text{Br} > \text{CH}_3 - \text{I}$   
 (D)  $\text{CH}_3 - \text{Cl} > \text{CH}_3 - \text{F} > \text{CH}_3 - \text{Br} > \text{CH}_3 - \text{I}$

Ans. (B)

Sol. The size of halogen atom increases from F to I hence bond length from C - F to C - I increases  $\therefore$  Bond enthalpy from  $\text{CH}_3 - \text{F}$  to  $\text{CH}_3 - \text{I}$  decreases

C - X Bond	Bond dissociation enthalpies/ $\text{kJ mol}^{-1}$
$\text{CH}_3 - \text{F}$	452
$\text{CH}_3 - \text{Cl}$	351
$\text{CH}_3 - \text{Br}$	293
$\text{CH}_3 - \text{I}$	234

69. Right option for the number of tetrahedral and octahedral voids in hexagonal primitive unit cell are :

- (A) 8, 4                      (B) 6, 12                      (C) 2, 1                      (D) 12, 6

Ans. (D)

Sol. • Number of octahedral and tetrahedral voids formed by N closed packed atoms are N and 2N respectively.

• Each hexagonal unit cell contains 6 atoms therefore, number of tetrahedral and octahedral voids are 12 and 6 respectively.

70. Which of the following reactions is the metal displacement reaction? Choose the right option.


- (A)  $2\text{KClO}_3 \xrightarrow{\Delta} 2\text{KCl} + 3\text{O}_2$   
 (B)  $\text{Cr}_2\text{O}_3 + 2\text{Al} \xrightarrow{\Delta} \text{Al}_2\text{O}_3 + 2\text{Cr}$   
 (C)  $\text{Fe} + 2\text{HCl} \rightarrow \text{FeCl}_2 + \text{H}_2 \uparrow$   
 (D)  $2\text{Pb}(\text{NO}_3)_2 \rightarrow 2\text{PbO} + 4\text{NO}_2 + \text{O}_2 \uparrow$

Ans. (B)

Sol. • Both reactions (A) and (D) are examples of decomposition reactions.

• Reactions (B) and (C), both are examples of displacement reactions, while reaction (B) is an example of metal displacement reaction.

71. Choose the correct option for graphical representation of Boyle's law, which shows a graph of pressure vs. volume of a gas at different temperatures :



Ans. (D)

Sol. According to Boyle's law

$$P \propto \frac{1}{V} \Rightarrow P = \frac{k}{V} \Rightarrow PV = k$$

where  $k$  is proportionality constant and equal to  $nRT$ .

$\therefore$  Graph between  $P$  vs.  $V$  should be rectangular hyperbola and product of  $PV$  increases with increase in temperature.

72. The  $pK_b$  of dimethylamine and  $pK_a$  of acetic acid are 3.27 and 4.77 respectively at  $T$  (K). The correct option for the pH of dimethylammonium acetate solution is :

(A) 8.50                      (B) 5.50                      (C) 7.75                      (D) 6.25

Ans. (C)

Sol. Dimethylammonium acetate is a salt of weak acid and weak base whose pH can be calculated as

$$\text{pH} = 7 + \frac{1}{2}(\text{p}K_a - \text{p}K_b)$$

$$= 7 + \frac{1}{2}(4.77 - 3.27)$$

$$= 7.75$$

73. Among the following alkaline earth metal halides, one which is covalent and soluble in organic solvents is :

(A) Calcium chloride  
 (B) Strontium chloride  
 (C) Magnesium chloride  
 (D) Beryllium chloride

Ans. (D)

Sol. • Except for beryllium chloride all other chloride of alkaline earth metals are ionic in nature.  
 • Due to small size of Be, Beryllium chloride is essentially covalent and soluble in organic solvents.

74. The maximum temperature that can be achieved in blast furnace is :
- (A) Upto 1200 K  
 (B) Upto 2200 K  
 (C) Upto 1900 K  
 (D) Upto 5000 K


**Ans. (B)**

**Sol.** Maximum temperature that can be achieved in blast furnace is upto 2200 K.  
 (As per NCERT text: 2170 K maximum temperature is given in the figure of blast furnace)

75. Ethylenediamine tetraacetate (EDTA) ion is :
- (A) Hexadentate ligand with four "O" and two "N" donor atoms  
 (B) Unidentate ligand  
 (C) Bidentate ligand with two "N" donor atoms  
 (D) Tridentate ligand with three "N" donor atoms

**Ans. (A)**

**Sol.** Ethylenediamine tetraacetate (EDTA) ion is a hexadentate ligand having four donor oxygen atoms and two donor nitrogen atoms



76. The following solutions were prepared by dissolving 10 g of glucose ( $C_6H_{12}O_6$ ) in 250 ml of water ( $P_1$ ), 10 g of urea ( $CH_4N_2O$ ) in 250 ml of water ( $P_2$ ) and 10 g of sucrose ( $C_{12}H_{22}O_{11}$ ) in 250 ml of water ( $P_3$ ).

The right option for the decreasing order of osmotic pressure of these solutions is :

- (A)  $P_2 > P_1 > P_3$   
 (B)  $P_1 > P_2 > P_3$   
 (C)  $P_2 > P_3 > P_1$   
 (D)  $P_3 > P_1 > P_2$

**Ans. (A)**

**Sol.** • Osmotic pressure ( $\pi$ ) =  $iCRT$

where C is molar concentration of the solution

- With increase in molar concentration of solution osmotic pressure increases.
- Since, weight of all solutes and its solution volume are equal, so higher will be the molar mass of solute, smaller will be molar concentration and smaller will be the osmotic pressure.
- Order of molar mass of solute decreases as Sucrose > Glucose > Urea
- So, correct order of osmotic pressure of solution is  $P_3 < P_1 < P_2$

77. **Statement I** : Acid strength increases in the order given as  $\text{HF} \ll \text{HCl} \ll \text{HBr} \ll \text{HI}$ .

**Statement II** : As the size of the elements F, Cl, Br, I increases down the group, the bond strength of HF, HCl, HBr and HI decreases and so the acid strength increases.


In the light of the above statements, choose the correct answer from the options given below.

- (A) Both statement I and Statement II are true
- (B) Both Statement I and Statement II are false
- (C) Statement I is correct but statement II is false
- (D) Statement I is incorrect but Statement II is true

**Ans.** (A)

**Sol.** In the modern periodic table, moving down the group as the size of halogen atom increases, the H - X bond length also increases as a result the bond enthalpy decreases. Hence, The acidic strength also increases.

So, the correct order of acidic strength is




78. The structures of beryllium chloride in solid state and vapour phase, are :


- (A) Chain and dimer, respectively
- (B) Linear in both
- (C) Dimer and Linear, respectively
- (D) Chain in both

**Ans.** (A)


**Sol.** Beryllium chloride has a chain structure in the solid state as shown below



In vapour phase Beryllium chloride tends to form a chloro-bridged dimer



79. For a reaction  $A \rightarrow B$ , enthalpy of reaction is  $-4.2 \text{ kJ mol}^{-1}$  and enthalpy of activation is  $9.6 \text{ kJ mol}^{-1}$ . The correct potential energy profile for the reaction is shown in option.



Ans. (B)

Sol. •  $\Delta H_{\text{rxn}} = (E_a)_f - (E_a)_b$

$$-4.2 = (E_a)_f - (E_a)_b$$

$$-4.2 = 9.6 - (E_a)_b$$

$$(E_a)_b = 9.6 + 4.2 = 13.8 \text{ kJ mol}^{-1}$$

- Since reaction is exothermic, so possible graph is (B) only.
- Also  $(E_a)_f < (E_a)_b$ , so answer is option (B).

80. Zr ( $Z = 40$ ) and Hf ( $Z = 72$ ) have similar atomic and ionic radii because of :

- (A) Belonging to same group  
 (B) Diagonal relationship  
 (C) Lanthanoid contraction  
 (D) Having similar chemical properties

Ans. (C)

- Sol. • The cumulative effect of the contraction of the lanthanoid series, known as lanthanoid contraction, causes the radii of the members of the third transition series to be very similar to those of the corresponding members of the second series.  
 • The almost identical radii of Zr (160 pm) and Hf (159 pm) is a consequence of the lanthanoid contraction.

81. A particular station of All India Radio, New Delhi broadcasts on a frequency of 1,368 kHz (kilohertz). The wavelength of the electromagnetic radiation emitted by the transmitter is : [speed of light  $c = 3.0 \times 10^8 \text{ ms}^{-1}$ ]

- (A) 219.3 m                      (B) 219.2 m                      (C) 2192 m                      (D) 21.92 cm

Ans. (A)

Sol. Energy of electromagnetic radiation (E)

$$= \frac{hc}{\lambda} = h\gamma$$

$$\text{So, } \frac{c}{\lambda} = \gamma \Rightarrow \lambda = \frac{c}{\gamma}$$

$$\lambda = \frac{3 \times 10^8}{1368 \times 10^3} = 219.3 \text{ m}$$

82. An organic compound contains 78% (by wt.) carbon and remaining percentage of hydrogen. The right option for the empirical formula of this compound is : [Atomic wt. of C is 12, H is 1]  
 (A) CH (B) CH<sub>2</sub> (C) CH<sub>3</sub> (D) CH<sub>4</sub>

Ans. (C)

Sol.


Element	Mass percentage	No. of mole	Mole ratio
C	78%	$\frac{78}{12} = 6.5$	$\frac{6.5}{6.5} = 1$
H	22%	$\frac{22}{1} = 22$	$\frac{22}{6.5} = 3.38 \approx 3$

Based on above calculation, possible empirical formula is CH<sub>3</sub>.

83. The compound which shows metamerism is :  
 (A) C<sub>5</sub>H<sub>12</sub> (B) C<sub>3</sub>H<sub>8</sub>O (C) C<sub>3</sub>H<sub>6</sub>O (D) C<sub>4</sub>H<sub>10</sub>O


Ans. (D)

Sol. Compounds with formula C<sub>4</sub>H<sub>10</sub>O can be ethers which may exhibit metamerism.  
 For example




and CH<sub>3</sub> - O - CH<sub>2</sub> - CH<sub>2</sub> - CH<sub>3</sub> are metamers as structure of alkyl chains are different around the functional group.

84. Identify the compound that will react with Hinsberg's reagent to give a solid which dissolves in alkali.




Ans. (C)

Sol. • Benzenesulphonyl chloride (C<sub>6</sub>H<sub>5</sub>SO<sub>2</sub>Cl) is also known as Hinsberg's reagent.  
 • The reaction of Hinsberg's reagent (C<sub>6</sub>H<sub>5</sub>SO<sub>2</sub>Cl) with primary amine (CH<sub>3</sub>CH<sub>2</sub>NH<sub>2</sub>) yields N-ethylbenzene sulphonamide.



• The reaction of Hinsberg's reagent (C<sub>6</sub>H<sub>5</sub>SO<sub>2</sub>Cl) with secondary amine (C<sub>2</sub>H<sub>5</sub>NHCH<sub>3</sub>) gives, N-Ethyl-N-Methyl benzene sulphonamide



• 3° amine do not react with Hinsberg reagent

85. The correct option for the number of body centred unit cells in all 14 types of Bravais lattice unit cells is :

- (A) 7 (B) 5 (C) 2 (D) 3

Ans. (D)

Sol. • In 14 types of Bravais lattices, body centred unit cell is present in cubic, tetragonal and orthorhombic crystal systems.

• Hence, body centred possible variation is present in three crystal systems.

### SECTION-B

86. Match List-I with List-II.

List-I	List-II
(a) $[\text{Fe}(\text{CN})_6]^{3-}$	(i) 5.92 BM
(b) $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$	(ii) 0 BM
(c) $[\text{Fe}(\text{CN})_6]^{4-}$	(iii) 4.90 BM
(d) $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$	(iv) 1.73 BM

Choose the correct answer from the options given below.

- (A) (a)-(iv), (b)-(ii), (c)-(i), (d)-(iii)  
 (B) (a)-(ii), (b)-(iv), (c)-(iii), (d)-(i)  
 (C) (a)-(i), (b)-(iii), (c)-(iv), (d)-(ii)  
 (D) (a)-(iv), (b)-(i), (c)-(ii), (d)-(iii)

Ans. (D)

Sol. Magnetic moment,  $\mu = \sqrt{n(n+2)}\text{BM}$

n = number of unpaired electrons)

Complex	No. of unpaired electron(s)	$\mu(\text{BM})$
(A) $[\text{Fe}(\text{CN})_6]^{3-}$	1	1.73
(B) $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$	5	5.92
(C) $[\text{Fe}(\text{CN})_6]^{4-}$	0	0
(D) $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$	4	4.90

87. Choose the correct option for the total pressure (in atm.) in a mixture of 4 g  $\text{O}_2$  and 2 g  $\text{H}_2$  confined in a total volume of one litre at  $0^\circ\text{C}$  is :

[Given  $R = 0.082 \text{ L atm mol}^{-1}\text{K}^{-1}$ ,  $T = 273 \text{ K}$ ]

- (A) 2.518 (B) 2.602 (C) 25.18 (D) 26.02

Ans. (C)


Sol.  $n_{\text{O}_2} = \frac{4}{32} = \frac{1}{8}$

$$n_{\text{H}_2} = \frac{2}{2} = 1$$

$$n_t = \frac{1}{8} + 1 = \frac{9}{8}$$

$$P_t V = n_t RT$$

$$P_t = \frac{\frac{9}{8} \times 0.082 \times 273}{1} = 25.18 \text{ atm}$$




Consider the above reaction and identify the missing reagent/chemical.

- (A)  $\text{B}_2\text{H}_6$                       (B) Red Phosphorus    (C) CaO                      (D) DIBAL-H

**Ans.** (C)

**Sol.** Alkane is produced by heating sodium salt of carboxylic acid with sodalime (NaOH and CaO in the ratio of 3 : 1)



89. For irreversible expansion of an ideal gas under isothermal condition, the correct option is:

- (A)  $\Delta U = 0, \Delta S_{\text{total}} = 0$                       (B)  $\Delta U \neq 0, \Delta S_{\text{total}} \neq 0$   
 (C)  $\Delta U = 0, \Delta S_{\text{total}} \neq 0$                       (D)  $\Delta U \neq 0, \Delta S_{\text{total}} = 0$

**Ans.** (C)

**Sol.** • For a spontaneous process,  $\Delta S_{\text{total}} > 0$  and since irreversible process is always spontaneous therefore  $\Delta S_{\text{total}} > 0$ .

• Since  $\Delta U = nC_V\Delta T$  and  $\Delta T = 0$  for isothermal process therefore  $\Delta U = 0$ .

90. In which one of the following arrangements the given sequence is not strictly according to the properties indicated against it?

- (A)  $\text{HF} < \text{HCl} < \text{HBr} < \text{HI}$  : Increasing acidic strength  
 (B)  $\text{H}_2\text{O} < \text{H}_2\text{S} < \text{H}_2\text{Se} < \text{H}_2\text{Te}$  : Increasing  $\text{pK}_a$  values  
 (C)  $\text{NH}_3 < \text{PH}_3 < \text{AsH}_3 < \text{SbH}_3$  : Increasing acidic character  
 (D)  $\text{CO}_2 < \text{SiO}_2 < \text{SnO}_2 < \text{PbO}_2$  : Increasing oxidizing power

**Ans.** (B)

**Sol.** Stronger is the acid, lower is the value of  $\text{pK}_a$ . On moving down the group, bond dissociation enthalpy of hydrides of group 16 elements decreases hence acidity increases and  $\text{pK}_a$  value decreases. Correct order of  $\text{pK}_a$  value will be  $\text{H}_2\text{O} > \text{H}_2\text{S} > \text{H}_2\text{Se} > \text{H}_2\text{Te}$

91. The molar conductivity of 0.007 M acetic acid is  $20 \text{ Scm}^2\text{mol}^{-1}$ . What is the dissociation constant of acetic acid? Choose the correct option.

$$\left[ \begin{array}{l} \Lambda_{\text{H}^+}^\circ = 350 \text{ Scm}^2 \text{ mol}^{-1} \\ \Lambda_{\text{CH}_3\text{COO}^-}^\circ = 50 \text{ Scm}^2 \text{ mol}^{-1} \end{array} \right]$$

- (A)  $1.75 \times 10^{-4} \text{ mol L}^{-1}$                       (B)  $2.50 \times 10^{-4} \text{ mol L}^{-1}$   
 (C)  $1.75 \times 10^{-5} \text{ mol L}^{-1}$                       (D)  $2.50 \times 10^{-5} \text{ mol L}^{-1}$

**Ans.** (C)

**Sol.**  $\Lambda_m = 20 \text{ Scm}^2 \text{ mol}^{-1}$

$$\Lambda_{\text{mCH}_3\text{COOH}}^\circ = \Lambda_{\text{CH}_3\text{COO}^-}^\circ + \Lambda_{\text{mH}^+}^\circ$$

$$= 50 + 350 = 400 \text{ Scm}^2 \text{ mol}^{-1}$$

$$\alpha = \frac{\Lambda_m}{\Lambda_m^\circ} = \frac{20}{400} = \frac{1}{20}$$

$$K_a = \frac{C\alpha^2}{1-\alpha} \approx C\alpha^2 = 7 \times 10^{-3} \times \left(\frac{1}{20}\right)^2$$

$$= 7 \times 10^{-3} \times \frac{1}{4} \times 10^{-2}$$

$$= 1.75 \times 10^{-5} \text{ mol L}^{-1}$$

92. The slope of Arrhenius plot  $\left(\ln k \text{ v/s } \frac{1}{T}\right)$  of first order reaction is  $-5 \times 10^3 \text{K}$ . The value of  $E_a$  of the reaction is. Choose the correct option for your answer.

[Given  $R = 8.314 \text{ JK}^{-1}\text{mol}^{-1}$ ]

- (A)  $41.5 \text{ kJ mol}^{-1}$  (B)  $83.0 \text{ kJ mol}^{-1}$   
 (C)  $166 \text{ kJ mol}^{-1}$  (D)  $-83 \text{ kJ mol}^{-1}$

Ans. (A)

Sol. Arrhenius equation

$$k = Ae^{-E_a/RT}$$

$$\ln k = \ln A + \ln e^{-E_a/RT}$$

$$\ln k = \ln A - \frac{E_a}{R} \left(\frac{1}{T}\right)$$

Slope of  $\ln k$  vs  $\frac{1}{T}$  curve,

$$m = -\frac{E_a}{R}$$






$$-5 \times 10^3 = -\frac{E_a}{R}$$

$$E_a = 5 \times 10^3 \times 8.314 \text{ J/mol}$$

$$= 41.57 \times 10^3 \text{ J/mol}$$


$$\approx 41.5 \text{ kJ/mol}$$

93. The product formed in the following chemical reaction is:







Ans. (D)

Sol.  $\text{NaBH}_4$  is a reducing agent. It reduces carbonyl group into alcohols but does not reduce esters.



94. Match List-I with List-II

List-I



List-II

(i) Hell-Volhard-Zelinsky reaction

(ii) Gattermann-Koch reaction

(iii) Haloform reaction

(iv) Esterification

Choose the correct answer from the options given below.

(A) (a) - (iv), (b) - (i), (c) - (ii), (d) - (iii)

(B) (a) - (iii), (b) - (ii), (c) - (i), (d) - (iv)


(C) (a) - (i), (b) - (iv), (c) - (iii), (d) - (ii)

(D) (a) - (ii), (b) - (iii), (c) - (iv), (d) - (i)

Ans. (D)

Sol.


Gattermann-Koch reaction:




Haloform reaction:



Esterification:




Hell-Volhard-Zelinsky reaction:






95. Which of the following molecules is non-polar in nature?  
 (A)  $\text{POCl}_3$  (B)  $\text{CH}_2\text{O}$  (C)  $\text{SbCl}_5$  (D)  $\text{NO}_2$

Ans. (C)

Sol.  $\text{SbCl}_5$  :



Net vector summation of bond moments will be zero so  $\text{SbCl}_5$  is a non-polar molecule.



96. From the following pairs of ions which one is not an iso-electronic pair?  
 (A)  $\text{O}^{2-}, \text{F}^-$  (B)  $\text{Na}^+, \text{Mg}^{2+}$  (C)  $\text{Mn}^{2+}, \text{Fe}^{3+}$  (D)  $\text{Fe}^{2+}, \text{Mn}^{2+}$

Ans. (D)

Sol. • Isoelectronic species have same number of electrons.

Species	Number of electrons
$\text{Fe}^{2+}$	$26 - 2 = 24$
$\text{Mn}^{2+}$	$25 - 2 = 23$
$\text{O}^{2-}$	$8 + 2 = 10$
$\text{F}^-$	$9 + 1 = 10$
$\text{Na}^+$	$11 - 1 = 10$
$\text{Mg}^{2+}$	$12 - 2 = 10$
$\text{Fe}^{3+}$	$26 - 3 = 23$

97. The correct option for the value of vapour pressure of a solution at  $45^\circ\text{C}$  with benzene to octane in molar ratio 3 : 2 is :

[At  $45^\circ\text{C}$  vapour pressure of benzene is 280 mm Hg and that of octane is 420 mm Hg. Assume Ideal gas]

- (A) 160 mm of Hg (B) 168 mm of Hg (C) 336 mm of Hg (D) 350 mm of Hg

Ans. (C)

Sol. Given :  $n_{\text{C}_6\text{H}_6} : n_{\text{C}_8\text{H}_{18}} = 3 : 2$

$$\text{So, } \chi_{\text{C}_6\text{H}_6} = \frac{3}{5}, \chi_{\text{C}_8\text{H}_{18}} = \frac{2}{5}$$




$$P_s = P_{\text{C}_6\text{H}_6}^{\circ} \chi_{\text{C}_6\text{H}_6} + P_{\text{C}_8\text{H}_{18}}^{\circ} \chi_{\text{C}_8\text{H}_{18}}$$

$$= 280 \times \frac{3}{5} + 420 \times \frac{2}{5}$$

$$= 168 + 168$$

$$= 336 \text{ mm of Hg}$$

98. Match List-I with List-II.

**List-I****List-II**

(i) Acid rain

(ii) Smog

(iii) Ozone depletion

(iv) Tropospheric pollution

Choose the correct answer from the options given below.

(A) (a)-(i), (b)-(ii), (c)-(iii), (d)-(iv)

(B) (a)-(ii), (b)-(iii), (c)-(iv), (d)-(i)

(C) (a)-(iv), (b)-(iii), (c)-(i), (d)-(ii)


(D) (a)-(iii), (b)-(ii), (c)-(iv), (d)-(i)

**Ans.** (C)

**Sol.** • Tropospheric pollution: In the presence of pollutant,  $\text{SO}_2$  converts into  $\text{SO}_3$ .



• In spring season, sunlight breaks HOCl and  $\text{Cl}_2$  to give chlorine radicals.




These chlorine radicals deplete ozone layer


• High level of sulphur causes acid rain which reacts with marble and causes discolouring and disfiguring



• A chain reaction occurs from interaction of NO with sunlight in which NO is converted to  $\text{NO}_2$  which absorb energy from sunlight and breaks into NO and O, which causes photochemical smog.



99. The reagent 'R' in the given sequence of chemical reaction is:



(A)  $\text{H}_2\text{O}$


(B)  $\text{CH}_3\text{CH}_2\text{OH}$

(C) HI

(D)  $\text{CuCN}/\text{KCN}$






**Ans.** (B)

**Sol.**




Reagent R  $\text{C}_2\text{H}_5\text{OH}$  with diazonium salt.

100. The intermediate compound 'X' in the following chemical reaction is:



Ans. (A)

Sol. Etard's reaction



**BOTANY (SECTION-A)**

- 101.** In spite of interspecific competition in nature, which mechanism the competing species might have evolved for their survival?  
 (A) Resource partitioning (B) Competitive release  
 (C) Mutualism (D) Predation

**Ans. (A)**

**Sol.** In spite of interspecific competition the competing species may co-exist by doing resource partitioning.

In mutualism two organisms are equally benefitted.

In predation one organism (Predator) eats the another one (Prey).

In competition release there occurs dramatical increase in population of a less distributed species when its superior competitor is removed.

- 102.** Match List-I with List-II.

	List - I		List - II
(a)	Cells with active cell division capacity	(i)	Vascular tissues
(b)	Tissue having all cells similar in structure and function	(ii)	Meristematic tissue
(c)	Tissue having different types of cells	(iii)	Sclereids
(d)	Dead cells with highly thickened walls and narrow lumen	(iv)	Simple tissue

Select the correct answer from the options given

- |           |       |       |       |
|-----------|-------|-------|-------|
| (a)       | (b)   | (c)   | (d)   |
| (A) (ii)  | (iv)  | (i)   | (iii) |
| (B) (iv)  | (iii) | (ii)  | (i)   |
| (C) (i)   | (ii)  | (iii) | (iv)  |
| (D) (iii) | (ii)  | (iv)  | (i)   |

**Ans. (A)**

- Sol.** (a) Meristematic tissues are those tissues which have cells with active cell division capacity.  
 (b) Simple tissues are those tissues which have all the cells similar in structure and function.  
 (c) Vascular tissues are complex permanent tissues hence they have different types of cells.  
 (d) Sclereids are sclerenchymatous cells which are dead with highly thickened walls and narrow lumen.

- 103.** During the purification process for recombinant DNA technology, addition of chilled ethanol precipitates out :  
 (A) RNA (B) DNA  
 (C) Histones (D) Polysaccharides

**Ans. (B)**

**Sol.** Various enzymes like protease, RNase, etc. are added to break down substances like proteins, RNA, etc. Once all these substances are broken down, DNA is left which is precipitated out by adding chilled ethanol.

Histones are basic proteins that help condense DNA in a cell.

104. Match List-I with List-II.

List-I		List-II	
(a)	Cohesion	(i)	More attraction in liquid phase
(b)	Adhesion	(ii)	Mutual attraction among water molecules
(c)	Surface tension	(iii)	Water loss in liquid phase
(d)	Guttation	(iv)	Attraction towards polar surfaces

Choose the correct answer from the options given below.

- |           |       |      |       |
|-----------|-------|------|-------|
| (a)       | (b)   | (c)  | (d)   |
| (1) (ii)  | (iv)  | (i)  | (iii) |
| (2) (iv)  | (iii) | (ii) | (i)   |
| (3) (iii) | (i)   | (iv) | (ii)  |
| (4) (ii)  | (i)   | (iv) | (iii) |

Ans. (A)

- Sol. (a) Cohesion is mutual attraction among water molecules.  
 (b) Adhesion is attraction towards polar surfaces.  
 (c) Surface tension explains water molecules are more attracted in liquid phase than gaseous phase.  
 (d) Guttation is loss of water in liquid form from the leaf margins.

105. The term used for transfer of pollen grains from anthers of one plant to stigma of a different plant which, during pollination, brings genetically different types of pollen grains to stigma, is :

- |                |                 |
|----------------|-----------------|
| (A) Xenogamy   | (B) Geitonogamy |
| (C) Chasmogamy | (D) Cleistogamy |

Ans. (A)

- Sol. Xenogamy refers to the transfer of pollen grains from anthers of one plant to stigma of a different plant which during pollination, brings genetically different types of pollen grains to stigma. Cleistogamy is a condition in which flower does not open. Geitonogamy refers to the transfer of pollen grain from anther to stigma of another flower of the same plant. Chasmogamy is a condition in which flowers remain open.

106. Which of the following stages of meiosis involves division of centromere?

- |                 |                  |                 |                  |
|-----------------|------------------|-----------------|------------------|
| (A) Metaphase I | (B) Metaphase II | (C) Anaphase II | (D) Telophase II |
|-----------------|------------------|-----------------|------------------|

Ans. (C)

Sol. Division of centromere occurs in anaphase II.

Telophase II is the last stage of meiosis II.

During this phase, the chromatids reach the poles and start uncoiling.

Chromosomes form two parallel plates in metaphase I and one plate in metaphase II.

107. Which of the following is a correct sequence of steps in a PCR (Polymerase Chain Reaction)?
- (A) Denaturation, Annealing, Extension  
 (B) Denaturation, Extension, Annealing  
 (C) Extension, Denaturation, Annealing  
 (D) Annealing, Denaturation, Extension

**Ans. (A)**

**Sol.** The first step in the polymerase chain reaction is denaturation during which strands of ds DNA separate. This requires temperature around 94°C. This is followed by annealing in which primers Annealing to 3' end of template DNA strand. Annealing is followed by extension in which Taq polymerase adds nucleotides to 3'OH end of primers.

108. Gemmae are present in
- (A) Mosses  
 (B) Pteridophytes  
 (C) Some Gymnosperms  
 (D) Some Liverworts

**Ans. (D)**

**Sol.** Gemmae are green, multicellular asexual buds that are produced by some liverworts like Marchantia. Mosses reproduce vegetatively by fragmentation and budding of protonema. Pteridophytes and Gymnosperms normally do not reproduce asexually

109. The production of gametes by the parents, formation of zygotes, the  $F_1$  and  $F_2$  plants, can be understood from a diagram called :
- (A) Bullet square  
 (B) Punch square  
 (C) Punnett square  
 (D) Net square

**Ans. (C)**

**Sol.** The production of gametes ( $n$ ) by the parents ( $2n$ ), the formation of the zygote ( $2n$ ), the  $F_1$  and  $F_2$  plants can be understood from a diagram called Punnett square.

110. The factor that leads to Founder effect in a population is :
- (A) Natural selection  
 (B) Genetic recombination  
 (C) Mutation  
 (D) Genetic drift

**Ans. (D)**

**Sol.** Change in gene frequency in a small population by chance is known as genetic drift. Genetic drift has two ramifications, one is bottle neck effect and another is founder's effect. When accidentally a few individuals are dispersed and act as founders of a new isolated population, founder's effect is said to be observed. Crossing over which occurs during gamete formation results in genetic recombination. Mutations are random and directionless.

111. Genera like Selaginella and Salvinia produce two kinds of spores. Such plants are known as:  
 (A) Homosorus (B) Heterosorus  
 (C) Homosporous (D) Heterosporous

**Ans. (D)**

**Sol.** Plants like Selaginella and Salvinia produce two kinds of spore i.e., microspores and macrospores. They are known as heterosporous.

Most of the pteridophytes produce single type of spores and are called homosporous. Sorus are brownish or yellowish cluster of sporeproducing structures located on the lower surface of fern leaves.

112. Plants follow different pathways in response to environment or phases of life to form different kinds of structures. This ability is called  
 (A) Elasticity (B) Flexibility  
 (C) Plasticity (D) Maturity

**Ans. (C)**

**Sol.** Plants show plasticity which means the ability of plant to follow different pathways and produce different structures in response to environment.

113. Which of the following are not secondary metabolites in plants?  
 (A) Morphine, codeine (B) Amino acids, glucose  
 (C) Vinblastin, curcumin (D) Rubber, gums

**Ans. (B)**

**Sol.** The correct option is (B)

Amino acids and glucose are included under the category of primary metabolites as they have identifiable functions and play known roles in normal physiological processes.

Rubber, gums, morphine, codeine, vinblastin and curcumin are included under the category of secondary metabolites as their role or functions in host organisms is not known yet. However, many of them are useful to human welfare.

114. Complete the flow chart on central dogma.

(a)  $\text{DNA} \xrightarrow{(b)} \text{mRNA} \xrightarrow{(c)} (d)$

- (A) (a)-Replication; (b)-Transcription;  
 (c)-Transduction; (d)-Protein  
 (B) (a)-Translation; (b)-Replication;  
 (c)-Transcription; (d)-Transduction  
 (C) (a)-Replication; (b)-Transcription;  
 (c)-Translation; (d)-Protein  
 (D) (a)-Transduction; (b)-Translation;  
 (c)-Replication; (d)-Protein

**Ans. (C)**

**Sol.** Formation of DNA from DNA is replication.

Formation of mRNA from DNA is called Transcription.

Formation of protein from mRNA is called Translation.

So, (a) is Replication

(b) is Transcription

(c) is Translation

(d) is Protein

Transduction is transfer of genetic material from one bacterium to another with the help of virus or a bacteriophage.

**115.** When the centromere is situated in the middle of two equal arms of chromosomes, the chromosome is referred as :

- (A) Metacentric (B) Telocentric  
(C) Sub-metacentric (D) Acrocentric

**Ans. (A)**

**Sol.** When the centromere is situated in the middle of two equal arms of chromosomes, the chromosome is referred as Metacentric.

When the centromere is present slightly away from the middle, it is called sub-metacentric chromosome.

When the centromere is present very close to one end of the chromosome, it is called acrocentric chromosome.

When the centromere is present at terminal position, the chromosome is called telocentric.

**116.** DNA strands on a gel stained with ethidium bromide when viewed under UV radiation, appear as

- (A) Yellow bands (B) Bright orange bands  
(C) Dark red bands (D) Bright blue bands

**Ans. (B)**

**Sol.** After the bands are stained, they are viewed in UV light. The bands appear bright orange in colour. Ethidium bromide is the intercalating agent that stacks in between the nitrogenous bases.

**117.** The site of perception of light in plants during photoperiodism is

- (A) Shoot apex (B) Stem  
(C) Axillary bud (D) Leaf

**Ans. (D)**

**Sol.** The site of perception of light in plants during photoperiodism is leaf.

The site of perception of low temperature stimulus during vernalisation is shoot apex and embryo. Axillary bud are not sites of perception of photoperiod.

**118.** When gene targetting involving gene amplification is attempted in an individual's tissue to treat disease, it is known as :

- (A) Biopiracy (B) Gene therapy  
(C) Molecular diagnosis (D) Safety testing

**Ans. (B)**

**Sol.** The correct option is (B)

Gene therapy is a collection of methods that allows correction of a gene defect that has been diagnosed in a child/embryo.

Biopiracy is the term used to refer to the use of bio-resources by multinational companies and other organisations without proper authorisation from the countries and people concerned without compensatory payment.

Molecular diagnosis refers to the act or process of determining the nature and cause of a disease.

119. Which of the following plants is monoecious?  
 (A) *Carica papaya* (B) *Chara*  
 (C) *Marchantia polymorpha* (D) *Cycas circinalis*

**Ans. (B)**

**Sol.** When male and female sex organs are present on same plant body, such plants are said to be monoecious.

Most of the species of *Chara* are monoecious.

*Cycas circinalis*, *Carica papaya* and *Marchantia polymorpha* are dioecious.

120. Which of the following is not an application of PCR (Polymerase Chain Reaction)?  
 (A) Molecular diagnosis (B) Gene amplification  
 (C) Purification of isolated protein (D) Detection of gene mutation

**Ans. (C)**

**Sol.** PCR is Polymerase Chain Reaction.

It is used for making multiple copies of the gene.

Hence PCR is used for

Gene amplification.

PCR-based assays have been developed that detect the presence of gene sequences of the infectious agents.

It is also used in detecting mutations.

Protein is not the target of PCR. Hence, plays no role in its purification.

121. Match List-I with List-II.

List-I		List-II	
(a)	Cristae	(i)	Primary constriction in chromosome
(b)	Thylakoids	(ii)	Disc-shaped sacs in Golgi apparatus
(c)	Centromere	(iii)	Infoldings in mitochondria
(d)	Cisternae	(iv)	Flattened membranous sacs in stroma of plastids

Choose the correct answer from the options given below.

- |           |       |       |      |
|-----------|-------|-------|------|
| (a)       | (b)   | (c)   | (d)  |
| (A) (iv)  | (iii) | (ii)  | (i)  |
| (B) (i)   | (iv)  | (iii) | (ii) |
| (C) (iii) | (iv)  | (i)   | (ii) |
| (D) (ii)  | (iii) | (iv)  | (i)  |

**Ans. (C)**

**Sol.** The inner membrane of mitochondria forms infoldings called cristae.

Thylakoids are flattened membranous sacs in stroma of plastids.

Cisternae are disc shaped sacs in Golgi apparatus.

Primary constriction in chromosome that holds two chromatids together is called centromere.

Hence correct option is (C)- a(iii), b(iv), c(i), d(ii)

122. Diadelphous stamens are found in  
 (A) China rose (B) Citrus  
 (C) Pea (D) China rose and citrus

Ans. (C)

Sol. Stamens are said to be diadelphous when these are united in two bundles e.g. Pea. China rose has monadelphous stamens while, Citrus has polyadelphous stamens. Monadelphous stamens are grouped in single bundle whereas polyadelphous stamens occur in more than two bundles.

123. Match List-I with List-II

List-I		List-II	
(a)	Protoplast fusion	(i)	Totipotency
(b)	Plant tissue culture	(ii)	Pomato
(c)	Meristem culture	(iii)	Somaclones
(d)	Micropropagation	(iv)	Virus free plants

Choose the correct answer from the options given below.

- |           |       |      |       |
|-----------|-------|------|-------|
| (a)       | (b)   | (c)  | (d)   |
| (A) (iii) | (iv)  | (ii) | (i)   |
| (B) (ii)  | (i)   | (iv) | (iii) |
| (C) (iii) | (iv)  | (i)  | (ii)  |
| (D) (iv)  | (iii) | (ii) | (i)   |

Ans. (B)

Sol. Pomato is obtained as a result of protoplast fusion. Totipotency is a property of explant to develop into whole plant body during plant tissue culture. Virus free plants can be obtained through meristem culture. Somaclones are obtained by the process of micropropagation.

124. Amensalism can be represented as:  
 (A) Species A (-); Species B (0) (B) Species A (+); Species B (+)  
 (C) Species A (-); Species B (-) (D) Species A (+); Species B (0)

Ans. (A)

Sol. Amensalism is an interaction between two organisms of different species in which one species inhibits the growth of other species by secreting certain chemicals. The first species is neither get benefited nor harmed.

(+) : (0) interaction is observed in commensalism

(+) : (+) interaction is observed in mutualism.

(-) : (-) interaction is seen in competition

125. Which of the following is an incorrect statement?  
 (A) Mature sieve tube elements possess a conspicuous nucleus and usual cytoplasmic organelles  
 (B) Microbodies are present both in plant and animal cells  
 (C) The perinuclear space forms a barrier between the materials present inside the nucleus and that of the cytoplasm  
 (D) Nuclear pores act as passages for proteins and RNA molecules in both directions between nucleus and cytoplasm

Ans. (A)

Sol. A mature sieve tube elements possess a peripheral cytoplasm and a large central vacuole but lacks a nucleus. Rest of other statements are correct.

126. A typical angiosperm embryo sac at maturity is:

- (A) 8-nucleate and 7-celled
- (B) 7-nucleate and 8-celled
- (C) 7-nucleate and 7-celled
- (D) 8-nucleate and 8-celled

**Ans.** (A)

**Sol.** A typical angiospermic embryo sac has seven cells that are three antipodals, one central cell, one egg cell and two synergids.

The central cell has two polar nuclei, hence the embryo sac is eight nucleated.

127. Which of the following algae contains mannitol as reserve food material?

- (A) Ectocarpus
- (B) Gracilaria
- (C) Volvox
- (D) Ulothrix

**Ans.** (A)

**Sol.** Ectocarpus is a brown alga belongs to the class Phaeophyceae. Members of this class have mannitol and laminarin as stored food material.

Ulothrix and Volvox belong to Chlorophyceae (green algae). Members of this class have starch as reserve food material. Gracilaria is a member of red algae (Rhodophyceae). This class is characterised by having floridean starch as stored food material.

128. The plant hormone used to destroy weeds in a field

- (A) IAA
- (B) NAA
- (C) 2, 4-D
- (D) IBA

**Ans.** (C)

**Sol.** Some synthetic auxins are used as weedicides.

2,4-D is widely used to remove broad leaved weeds or dicotyledonous weeds in cereal crops or monocotyledonous plants.

IAA and IBA are natural auxins.

NAA is a synthetic auxin.

129. The amount of nutrients, such as carbon, nitrogen, phosphorus and calcium present in the soil at any given time, is referred as :

- (A) Climax
- (B) Climax community
- (C) Standing state
- (D) Standing crop

**Ans.** (C)

**Sol.** Amount of all the inorganic substances or nutrients, such as carbon, nitrogen, phosphorus and calcium present in soil at any given time, is referred as standing state.

Amount of living material present in different trophic levels at a given time, is referred as standing crop.

Climax community is the last community in biotic succession which is relatively stable and is in near equilibrium with the environment of that area.

130. Mutations in plant cells can be induced by:  
 (A) Kinetin (B) Infrared rays (C) Gamma rays (D) Zeatin

**Ans.** (C)

**Sol.** Several kinds of radiation like gamma rays, X-rays, UV-rays cause mutation.


These are physical mutagens.

Such induced mutation in plants is done to develop improved varieties. The first natural cytokinin was isolated from unripe maize grain known as zeatin. The cytokinin that was obtained from degraded product of autoclaved herring sperm DNA was kinetin (N<sup>6</sup>-furfuryl aminopurine). Infrared rays cause heating effect.

131. Which of the following statements is not correct?  
 (A) Pyramid of biomass in sea is generally inverted.  
 (B) Pyramid of biomass in sea is generally upright.  
 (C) Pyramid of energy is always upright.  
 (D) Pyramid of numbers in a grassland ecosystem is upright.

**Ans.** (B)

**Sol.** Pyramid of biomass in sea is inverted. For example, biomass of zooplanktons is higher than that of phytoplanktons as life span of former is longer and the latter multiply much faster though having shorter life span.



Small standing crop of phytoplankton supports large standing crop of zooplankton

132. In the equation  $GPP - R = NPP$   
 R represents :  
 (A) Radiant energy (B) Retardation factor  
 (C) Environmental factor (D) Respiration losses

**Ans.** (D)

**Sol.** In the equation,

$$GPP - R = NPP$$

R refers to respiratory loss

GPP is gross primary productivity

NPP is net primary productivity

133. Which of the following algae produce Carrageen?  
 (A) Green algae (B) Brown algae  
 (C) Red algae (D) Blue-green algae

**Ans.** (C)

**Sol.** The cell wall of red algae is composed of agar, carrageen along with cellulose.

In brown algae cell wall contains algin while in green algae it is composed of cellulose and pectin.

In blue green algae cell wall is composed of mucopeptides.

134. The first stable product of  $\text{CO}_2$  fixation in Sorghum is  
 (A) Pyruvic acid (B) Oxaloacetic acid  
 (C) Succinic acid (D) Phosphoglyceric acid

Ans. (B)

Sol. Sorghum is a  $\text{C}_4$  plant. The first stable product of  $\text{CO}_2$  fixation in Sorghum is oxaloacetic acid. The first stable product in  $\text{C}_3$  cycle is 3-phosphoglyceric acid. Pyruvic acid is the end product of glycolysis. Succinic acid is an intermediate product in krebs cycle.

135. Match List-I with List-II.

List-I		List-II	
(a)	Lenticels	(i)	Phellogen
(b)	Cork cambium	(ii)	Suberin deposition
(c)	Secondary cortex	(iii)	Exchange of gases
(d)	Cork	(iv)	Phelloderm

Choose the correct answer from the options given below.

- |           |       |       |       |
|-----------|-------|-------|-------|
| (a)       | (b)   | (c)   | (d)   |
| (A) (iv)  | (i)   | (iii) | (ii)  |
| (B) (iii) | (i)   | (iv)  | (ii)  |
| (C) (ii)  | (iii) | (iv)  | (i)   |
| (D) (iv)  | (ii)  | (i)   | (iii) |

Ans. (B)

Sol. Lenticels are meant for exchange of gases. Phellogen is also known as cork cambium. Phelloderm is also called secondary cortex because it is the cortex that develops during secondary growth. Cork has deposition of suberin in their cell walls when they get mature.

### SECTION - B

136. Which of the following statements of incorrect?  
 (A) During aerobic respiration, role of oxygen is limited to the terminal stage  
 (B) In ETC (Electron Transport Chain), one molecule of  $\text{NADH} + \text{H}^+$  gives rise to 2 ATP molecules, and one  $\text{FADH}_2$  gives rise to 3 ATP molecules  
 (C) ATP is synthesized through complex V  
 (D) Oxidation-reduction reactions produce proton gradient in respiration

Ans. (B)

Sol. During respiration, process of ATP synthesis is explained by chemiosmotic model. It says that a proton gradient is required for ATP synthesis that is established by oxidation-reduction reactions. In ETC, one  $\text{NADH} + \text{H}^+$  produces 3 ATP while one  $\text{FADH}_2$  produces 2 ATP molecules. ATP is synthesised via complex V. In ETS, oxygen acts as terminal electron acceptor.

## 137. Match Column-I with Column-II

Column-I	Column-II
(a) $\% \overline{\text{K}}_{(5)} \text{C}_{1+2+(2)} \text{A}_{(9)+1} \underline{\text{G}}_1$	(i) Brassicaceae
(b) $\oplus \overline{\text{K}}_{(5)} \overline{\text{C}}_{(5)} \overline{\text{A}}_5 \underline{\text{G}}_{(2)}$	(ii) Liliaceae
(c) $\oplus \overline{\text{P}}_{(3+3)} \overline{\text{A}}_{3+3} \underline{\text{G}}_{(3)}$	(iii) Fabaceae
(d) $\oplus \overline{\text{K}}_{2+2} \text{C}_4 \text{A}_{2-4} \underline{\text{G}}_{(2)}$	(iv) Solanaceae

Select the correct answer from the options given below.

(a)	(b)	(c)	(d)
(A) (iii)	(iv)	(ii)	(i)
(B) (i)	(ii)	(iii)	(iv)
(C) (ii)	(iii)	(iv)	(i)
(D) (iv)	(ii)	(i)	(iii)

Ans. (A)

Sol. The floral formula of

Brassicaceae family -  $\oplus \overline{\text{K}}_{2+2} \text{C}_4 \text{A}_{2+4} \underline{\text{G}}_{(2)}$

Solanaceae family -  $\oplus \overline{\text{K}}_{(5)} \overline{\text{C}}_{(5)} \overline{\text{A}}_5 \underline{\text{G}}_{(2)}$

Fabaceae family -  $\% \overline{\text{K}}_{(5)} \text{C}_{1+2+(2)} \text{A}_{(9)+1} \underline{\text{G}}_1$

Liliaceae family -  $\oplus \overline{\text{P}}_{(3+3)} \overline{\text{A}}_{3+3} \underline{\text{G}}_{(3)}$

So a(iii), b(iv), c(ii), d(i) is correct matching.

## 138. Match List-I with List-II.

List-I		List-II	
(a)	S phase	(i)	Proteins are synthesized
(b)	G <sub>2</sub> phase	(ii)	Inactive phase
(c)	Quiescent stage	(iii)	Interval between mitosis and initiation of DNA replication
(d)	G <sub>1</sub> phase	(iv)	DNA replication

Choose the correct answer from the options given below.

(a)	(b)	(c)	(d)
(A) (iii)	(ii)	(i)	(iv)
(B) (iv)	(ii)	(iii)	(i)
(C) (iv)	(i)	(ii)	(iii)
(4) (ii)	(iv)	(iii)	(i)

Ans. (C)

Sol. In S phase DNA replication takes place.

In G<sub>2</sub> phase there is synthesis of proteins, RNA etc.

Quiescent stage is inactive stage of cell cycle but cells remain metabolically active in this stage.

G<sub>1</sub> phase is the interval between mitosis and initiation of DNA replication.

139. Plasmid pBR322 has PstI restriction enzyme site within gene amp<sup>R</sup> that confers ampicillin resistance.

If this enzyme is used for inserting a gene for  $\beta$ -galactoside production and the recombinant plasmid is inserted in an E.coli strain

- (A) It will not be able to confer ampicillin resistance to the host cell
- (B) The transformed cells will have the ability to resist ampicillin as well as produce  $\beta$ -galactoside
- (C) It will lead to lysis of host cell
- (D) It will be able to produce a novel protein with dual ability

**Ans. (A)**

**Sol.** pBR322 is a commonly used cloning vector. When the gene for  $\beta$ -galactoside is inserted in the ampicillin resistance gene by using Pst I, the recombinant E.coli will lose ampicillin resistance due to insertional inactivation of the antibiotic resistance gene.

The host (recombinant) cell will produce  $\beta$ -galactoside which is not a novel protein nor does it have dual ability.

The transformed cells cannot resist ampicillin as they have lost ampicillin resistance.

A recombinant E. coli is produced and the host cell will not undergo lysis due to insertion of  $\beta$ -galactoside gene.

140. Identify the correct statement.

- (A) In capping, methyl guanosine triphosphate is added to the 3' end of hnRNA
- (B) RNA polymerase binds with Rho factor to terminate the process of transcription in bacteria
- (C) The coding strand in a transcription unit is copied to an mRNA
- (4) Split gene arrangement is characteristic of prokaryotes

**Ans. (B)**

**Sol.** Split gene arrangement is characteristic of eukaryotes.

In capping 5-methyl guanosine triphosphate is added at 5' end of hnRNA.

At 3' end poly-A tail is added.

The non coding or template strand is copied to an mRNA. RNA polymerase associate with  $\rho$  factor (Rho factor) and it alters the specificity of the RNA polymerase to terminate the processes.

141. Now a days it is possible to detect the mutated gene causing cancer by allowing radioactive probe to hybridise its complimentary DNA in a clone of cells, followed by its detection using autoradiography because :

- (A) Mutated gene partially appears on a photographic film
- (B) Mutated gene completely and clearly appears on a photographic film
- (C) Mutated gene does not appear on a photographic film as the probe has no complementarity with it
- (D) Mutated gene does not appear on photographic film as the probe has complementarity with it

**Ans. (C)**

**Sol.** Autoradiography allows the detection/localisation of radioactive isotope within a biological sample.

Probe is a radiolabelled ss DNA or ss RNA depending on the technique. To identify the mutated gene probe is allowed to hybridise to its complementary DNA in a clone of cells followed by detection using autoradiography. The mutated gene will not appear on the photographic film, because the probe does not have complementarity with the mutated gene.

- 142.** In the exponential growth equation  $N_t = N_0 e^{rt}$ ,  $e$  represents
- (A) The base of number logarithms  
 (B) The base of exponential logarithms  
 (C) The base of natural logarithms  
 (D) The base of geometric logarithms

**Ans.** (C)

**Sol.** In the exponential growth equation  $N_t = N_0 e^{rt}$ ,  $e$  represents the base of natural logarithms  
 $N_t$  = Population density after time  $t$   
 $N_0$  = Population density at time zero  
 $r$  = Intrinsic rate of natural increase called biotic potential.

- 143.** Select the correct pair.
- (A) Large colorless empty - Subsidiary cells cells in the epidermis of grass leaves  
 (B) In dicot leaves, vascular - Conjunctive bundles are surrounded tissue by large thick-walled cells  
 (C) Cells of medullary rays - Interfascicular that form part of cambium cambial ring  
 (4) Loose parenchyma cells - Spongy rupturing the epidermis parenchyma and forming a lens shaped opening in bark

**Ans.** (C)

**Sol.** When the cells of medullary rays differentiated, they give rise to the new cambium called interfascicular cambium.  
 Loose parenchyma cells rupturing the epidermis and forming a lens-shaped opening in bark are called complementary cells.  
 Large colourless empty cells in the epidermis of grass leaves are called bulliform cells.  
 In dicot leaf, vascular bundles are surrounded by large thick walled cells called bundle sheath cells.

- 144.** In some members of which of the following pairs of families, pollen grains retain their viability for months after release?
- (A) Poaceae ; Rosaceae  
 (B) Poaceae ; Leguminosae  
 (C) Poaceae ; Solanaceae  
 (D) Rosaceae ; Leguminosae

**Ans.** (D)

**Sol.** In members of some plant families like Solanaceae, Rosaceae and Leguminosae the pollen grains retain their viability for several months.  
 In cereals (Poaceae) pollen grains retain viability for around 30 minutes.

- 145.** What is the role of RNA polymerase III in the process of transcription in eukaryotes?
- (A) Transcribes rRNAs (28S, 18S and 5.8S)  
 (B) Transcribes tRNA, 5s rRNA and snRNA  
 (C) Transcribes precursor of mRNA  
 (D) Transcribes only snRNAs

**Ans.** (B)

**Sol.** RNA polymerase III transcribes tRNA, ScRNA, 5S rRNA and SnRNA.  
 RNA polymerase I transcribes 5.8S, 18S and 28S rRNA.  
 RNA polymerase II transcribes hnRNA which is precursor of mRNA

146. Which of the following statements is incorrect?
- (A) Both ATP and NADPH + H<sup>+</sup> are synthesized during non-cyclic photophosphorylation  
 (B) Stroma lamellae have PS I only and lack NADP reductase  
 (C) Grana lamellae have both PS I and PS II  
 (D) Cyclic photophosphorylation involves both PS I and PS II

Ans. (D)

Sol. Cyclic photophosphorylation involves only PS I.

Both PS I and PS II are involved in non-cyclic photophosphorylation where both ATP and NADPH + H<sup>+</sup> are synthesized.

Both PS I and PS II are found on grana lamellae whereas stroma lamellae have PS I only and lack NADP reductase.

147. Which of the following statements is correct ?
- (A) Fusion of two cells is called Karyogamy  
 (B) Fusion of protoplasts between two motile on non-motile gametes is called plasmogamy  
 (C) Organisms that depend on living plants are called saprophytes  
 (D) Some of the organisms can fix atmospheric nitrogen in specialized cells called sheath cells

Ans. (B)

Sol. In some blue-green algae specialised cells called heterocyst fixes atmospheric nitrogen into ammonia.

Fusion of two nuclei is called Karyogamy.

Organisms that depend on living plants are parasites, saprophytes grow on dead material.

Fusion of protoplasts of two cells is called plasmogamy.

148. Match List-I with List-II.

List-I		List-II
(a) Protein	(i)	C = C double bonds
(b) Unsaturated fatty acid	(ii)	Phosphodiester bonds
(c) Nucleic acid	(iii)	Glycosidic bonds
(d) Polysaccharide	(iv)	Peptide bonds

Choose the correct answer from the options given below.

- |          |       |       |       |
|----------|-------|-------|-------|
| (a)      | (b)   | (c)   | (d)   |
| (A) (iv) | (i)   | (ii)  | (iii) |
| (B) (i)  | (iv)  | (iii) | (ii)  |
| (C) (ii) | (i)   | (iv)  | (iii) |
| (D) (iv) | (iii) | (i)   | (ii)  |

Ans. (A)

Sol. In a polypeptide or a protein, amino acids are linked by a peptide bond which is formed when the carboxyl (-COOH) group of one amino acid reacts with amino (-NH<sub>2</sub>) group of the next amino acid with the elimination of a water moiety.

Unsaturated fatty acids are with one or more C = C double bonds.

In nucleic acids, a phosphate moiety links the 3'-carbon of one sugar of one nucleotide to the 5'-carbon of the sugar of the succeeding nucleotide. The bond between the phosphate and hydroxyl group is an ester bond. As there is one such ester bond on either side, it is called phosphodiester bond. In a polysaccharide, the individual monosaccharides are linked by a glycosidic bond.

149. DNA fingerprinting involves identifying differences in some specific regions in DNA sequence, called as
- (A) Satellite DNA (B) Repetitive DNA  
(C) Single nucleotides (D) Polymorphic DNA

**Ans. (B)**

**Sol.** DNA fingerprinting involves identifying differences in some specific regions in DNA sequence called as repetitive DNA.

The basis of DNA fingerprinting is VNTR (a satellite DNA as probe that show very high degree of polymorphism)

Polymorphism is the variation at genetic level.

Allelic sequence variation has traditionally been described as a DNA polymorphism.

150. Match Column-I with Column-II.

	Column-I		Column-II
(a)	<i>Nitrococcus</i>	(i)	Denitrification
(b)	<i>Rhizobium</i>	(ii)	Conversion of ammonia to nitrite
(c)	<i>Thiobacillus</i>	(iii)	Conversion of nitrite to nitrate
(d)	<i>Nitrobacter</i>	(iv)	Conversion of atmospheric nitrogen to ammonia

Choose the correct answer from options given below.

- |           |       |       |       |
|-----------|-------|-------|-------|
| (a)       | (b)   | (c)   | (d)   |
| (A) (ii)  | (iv)  | (i)   | (iii) |
| (B) (i)   | (ii)  | (iii) | (iv)  |
| (C) (iii) | (i)   | (iv)  | (ii)  |
| (D) (iv)  | (iii) | (ii)  | (i)   |

**Ans. (A)**

**Sol.** Nitrogen fixation is conversion of atmospheric  $N_2$  to  $NH_3$  (ammonia). It is carried out by  $N_2$  fixers such as *Rhizobium*.

$NH_3$  is converted to  $NO_2^-$  (nitrite) by nitrifying bacteria such as *Nitrococcus*.

Then is  $NO_2^-$  converted to  $NO_3^-$  (nitrate) by nitrifying bacteria called *Nitrobacter*.

*Thiobacillus* carries out denitrification, a process where  $NO_2^- / NO_3^-$  is converted to  $N_2$ .

## (ZOOLOGY) SECTION - A

**151.** A specific recognition sequence identified by endonucleases to make cuts at specific positions within the DNA is:

- (A) Degenerate primer sequence
- (B) Okazaki sequences
- (C) Palindromic Nucleotide sequences
- (D) Poly(A) tail sequences

**Ans.** (C)

**Sol.** Each restriction endonuclease recognizes a specific palindromic nucleotide sequence in the DNA. Once it finds its specific recognition sequence it binds to DNA and cuts each of the two strands of DNA.

During post transcriptional modification in eukaryotes, poly(A) tail (200–300 adenylate residues) are added at 3' end of hnRNA.

During DNA replication Okazaki fragments are synthesized discontinuously and joined by DNA ligase.

A PCR primer sequence is termed degenerate if some of its positions have several possible bases.

**152.** The fruit fly has 8 chromosomes (2n) in each cell.

During interphase of Mitosis if the number of chromosomes at G<sub>1</sub> phase is 8, what would be the number of chromosomes after S phase?

- (A) 8
- (B) 16
- (C) 4
- (D) 32

**Ans.** (A)

**Sol.** In S phase there is duplication of DNA. So amount of DNA increases but not the chromosome number.

So, if the number of chromosomes at G<sub>1</sub> phase is 8 in fruit fly then the number of chromosomes will be same in S phase that is 8 only.

**153.** Which one of the following belongs to the family Muscidae?

- (A) Fire fly
- (B) Grasshopper
- (C) Cockroach
- (D) House fly

**Ans.** (D)

**Sol.** Option (D) is correct because housefly belongs to the family Muscidae, class Insecta and phylum Arthropoda.

Fire flies are placed in family Lampyridae of class Insecta.

Grasshopper is also an insect placed in family Acrididae.

Cockroach is also an insect placed in family Blattidae.

**154.** Succus entericus is referred to as:

- (A) Pancreatic juice
- (B) Intestinal juice
- (C) Gastric juice
- (D) Chyme

**Ans.** (B)

**Sol.** Option (B) is correct because succus entericus is referred to as intestinal juice.

Chyme is the name given to acidic food present in stomach.

Exocrine secretion of pancreatic acini is called pancreatic juice.

Secretion of gastric glands present in stomach is called gastric juice.

155. With regard to insulin choose correct options.
- (a) C-peptide is not present in mature insulin.
  - (b) The insulin produced by rDNA technology has C-peptide.
  - (c) The pro-insulin has C-peptide
  - (d) A-peptide and B-peptide of insulin are interconnected by disulphide bridges.

Choose the correct answer from the options given below

- (A) (b) and (d) only
- (B) (b) and (c) only
- (C) (a), (c) and (d) only
- (D) (a) and (d) only

**Ans.** (C)

**Sol.** Insulin is synthesized as a pro-hormone which contains A-chain, B-chain and an extra stretch called the C-peptide.

C-peptide is not present in mature insulin called humulin.

Chains A and B are connected by interchain disulphide bridges.

156. Persons with 'AB' blood group are called as "Universal recipients". This is due to :

- (A) Absence of antigens A and B on the surface of RBCs
- (B) Absence of antigens A and B in plasma
- (C) Presence of antibodies, anti-A and anti-B, on RBCs
- (D) Absence of antibodies, anti-A and anti-B, in plasma

**Ans.** (D)


**Sol.** Option (D) is correct because persons with 'AB' blood group contain antigens 'A' and 'B' but lack antibodies anti-A and anti-B in plasma. So, persons with 'AB' blood group can accept blood from persons with AB as well as the other groups of blood due to lack of antibodies in their blood. Therefore, such persons are called "Universal recipients".

157. In a cross between a male and female, both heterozygous for sickle cell anaemia gene, what percentage of the progeny will be diseased?

- (A) 50%
- (B) 75%
- (C) 25%
- (D) 100%

**Ans.** (C)

**Sol.** According to given question;



Total number of affected progenies = 1

∴ Percentage of diseased/affected progenies

$$= \frac{1}{4} \times 100 = 25\%$$

158. Which enzyme is responsible for the conversion of inactive fibrinogens to fibrins?

- (A) Thrombin (B) Renin  
(C) Epinephrine (D) Thrombokinase

Ans. (A)

Sol. During coagulation of blood, an enzyme complex thrombokinase helps in the conversion of prothrombin (present in plasma) into thrombin.

Thrombin further helps in the conversion of inactive fibrinogens into fibrins which form network of threads.

Renin is secreted by JG cells in response to fall in glomerular blood flow, which converts angiotensinogen in blood to angiotensin-I

Epinephrine or adrenaline is secreted by adrenal medulla in response to stress of any kind and during emergency.

159. The partial pressures (in mm Hg) of oxygen ( $O_2$ ) and carbon dioxide ( $CO_2$ ) at alveoli (the site of diffusion) are:

- (A)  $pO_2 = 104$  and  $pCO_2 = 40$  (B)  $pO_2 = 40$  and  $pCO_2 = 45$   
(C)  $pO_2 = 95$  and  $pCO_2 = 40$  (D)  $pO_2 = 159$  and  $pCO_2 = 0.3$

Ans. (A)

Sol. Option (A) is correct because  $pO_2$  in alveoli is 104 mm Hg and  $pCO_2$  in alveoli is 40 mmHg.

In atmosphere,  $pO_2$  is 159 mm Hg and  $pCO_2$  is 0.3 mm Hg.

In deoxygenated blood,  $pO_2$  is 40 mmHg and  $pCO_2$  is 45 mmHg.

In oxygenated blood,  $pO_2$  is 95 mmHg and  $pCO_2$  is 40 mmHg.

160. Chronic auto immune disorder affecting neuro muscular junction leading to fatigue, weakening and paralysis of skeletal muscle is called as:

- (A) Arthritis (B) Muscular dystrophy  
(C) Myasthenia gravis (D) Gout

Ans. (C)

Sol. Option (C) is correct because myasthenia gravis is a chronic auto immune disorder affecting neuromuscular junction leading to fatigue, weakening and paralysis of skeletal muscle.

Gout is caused due to deposition of uric acid crystals in joints leading to its inflammation.

Inflammation of joints is commonly known as arthritis.

Muscular dystrophy is a genetic disorder which results in progressive degeneration of skeletal muscle.

161. Which is the "Only enzyme" that has "Capability" to catalyse Initiation, Elongation and Termination in the process of transcription in prokaryotes?

- (A) DNA dependent DNA polymerase (B) DNA dependent RNA polymerase  
(C) DNA Ligase (D) DNase

Ans. (B)

Sol. In prokaryotes, the DNA dependent RNA polymerase is a holoenzyme that is made of polypeptides ( $\alpha_2\beta\beta'\omega$ )  $\sigma$ . It is responsible for initiation, elongation and termination during transcription.

DNase degrades DNA.

DNA dependent DNA polymerase is involved in replication of DNA.

DNA ligase joins the discontinuously synthesised fragments of DNA.

162. Which of the following RNAs is not required for the synthesis of protein?

- (A) mRNA
- (B) tRNA
- (C) rRNA
- (D) siRNA

**Ans** (D)

**Sol.** siRNA are small interfering RNA also called silencing RNA. It is a class of double-stranded RNA, non-coding RNA molecules.

mRNA is messenger RNA that carries genetic information provided by DNA.

tRNA carries amino acids to the mRNA during translation.

rRNA is structural RNA that forms ribosomes which are involved in translation.

163. Which one of the following is an example of Hormone releasing IUD?

- (A) CuT
- (B) LNG 20
- (C) Cu 7
- (D) Multiload 375

**Ans.** (B)

**Sol.** LNG-20 is a hormone releasing IUD which makes the uterus unsuitable for implantation and the cervix hostile to sperms.

Multiload 375, CuT and Cu7 are copper releasing IUDs which suppress sperm motility and the fertilizing capacity of sperms.

164. If Adenine makes 30% of the DNA molecule, what will be the percentage of Thymine, Guanine and Cytosine in it?

- (A) T : 20 ; G : 30 ; C : 20
- (B) T : 20 ; G : 20 ; C : 30
- (C) T : 30 ; G : 20 ; C : 20
- (D) T : 20 ; G : 25 ; C : 25

**Ans.** (C)

**Sol.** According to Chargaff's rule, for a double stranded DNA,

$$[A] = [T],$$

$$\therefore [A] = 30\%, \Rightarrow [T] = 30\%$$

$$\text{Since } [C] = [G]$$

$$\therefore 100 - [A + T]$$

$$= 100 - [30 + 30]$$

$$= 100 - 60 = 40 \%$$

$$\text{and } C = G = 20\% \text{ each}$$

$$\therefore A 30\%$$

$$T 30\%$$

$$G 20\%$$

$$C 20\%$$

165. Match List-I with List-II

List-I		List-II	
(a)	<i>Aspergillus niger</i>	(i)	Acetic Acid
(b)	<i>Acetobacter aceti</i>	(ii)	Lactic Acid
(c)	<i>Clostridium butylicum</i>	(iii)	Citric Acid
(d)	<i>Lactobacillus</i>	(iv)	Butyric Acid

Choose the correct answer from the options given below

- |           |       |       |       |
|-----------|-------|-------|-------|
| (a)       | (b)   | (c)   | (d)   |
| (A) (iii) | (i)   | (iv)  | (ii)  |
| (B) (i)   | (ii)  | (iii) | (iv)  |
| (C) (ii)  | (iii) | (i)   | (iv)  |
| (D) (iv)  | (ii)  | (i)   | (iii) |

**Ans. (A)**

**Sol.** *Aspergillus niger* is involved in production of citric acid. *Acetobacter aceti* is involved in production of acetic acid. *Clostridium butylicum* is involved in production of butyric acid whereas *Lactobacillus* is involved in the production of lactic acid. So a(iii), b(i), c(iv), d(ii) is correct matching.

166. Read the following statements

- Metagenesis is observed in Helminths.
- Echinoderms are triploblastic and coelomate animals.
- Round worms have organ-system level of body organization.
- Comb plates present in ctenophores help in digestion.
- Water vascular system is characteristic of Echinoderms.

Choose the correct answer from the options given below.

- |                                  |                                  |
|----------------------------------|----------------------------------|
| (A) (c), (d) and (e) are correct | (B) (a), (b) and (c) are correct |
| (C) (a), (d) and (e) are correct | (D) (b), (c) and (e) are correct |

**Ans. (D)**

**Sol.** Metagenesis (alternation of generation) is observed in members of phylum Coelenterata (Cnidaria).

Echinoderms are triploblastic and coelomate animals as true coelom is observed in them.

Roundworms (Aschelminths) have organ system level of organization.

Comb plates present in ctenophores help in locomotion.

Water vascular system is seen in echinoderms, which helps in locomotion, capture and transport of food and respiration.

167. Receptors for sperm binding in mammals are present on :

- |                         |                        |
|-------------------------|------------------------|
| (A) Corona radiata      | (B) Vitelline membrane |
| (C) Perivitelline space | (D) Zona pellucida     |

**Ans. (D)**

**Sol.** Option (D) is correct because zona pellucida has receptors for sperm binding (ZP3 receptors) in mammals.

Corona radiata is a layer of radially arranged cells of membrana granulosa.

Perivitelline space is present in between vitelline membrane and zona pellucida.

## 168. Match List - I with List - II

List - I		List - II	
(a)	Metamerism	(i)	Coelenterata
(b)	Canal system	(ii)	Ctenophora
(c)	Comb plates	(iii)	Annelida
(d)	Cnidoblasts	(iv)	Porifera

Choose the correct answer from the options given below.

- |           |       |      |       |
|-----------|-------|------|-------|
| (a)       | (b)   | (c)  | (d)   |
| (A) (iv)  | (iii) | (i)  | (ii)  |
| (B) (iii) | (iv)  | (i)  | (ii)  |
| (C) (iii) | (iv)  | (ii) | (i)   |
| (D) (iv)  | (i)   | (ii) | (iii) |

**Ans. (C)**

**Sol.** Metamerism is commonly seen in the members of phylum Annelida where the body is externally and internally divided into segments with a serial repetition of atleast some organs.

Water canal system is present in the members of phylum Porifera.

The body of ctenophores bears 8 external rows of ciliated comb plates which help in locomotion. Cnidoblasts or cnidocytes are characteristic feature of cnidarians (coelenterata).

## 169. Erythropoietin hormone which stimulates R.B.C. formation is produced by:

- |                              |  |
|------------------------------|--|
| (A) Alpha cells of pancreas  | (B) The cells of rostral adenohypophysis |
| (C) The cells of bone marrow | (D) Juxtaglomerular cells of the kidney  |

**Ans. (D)**

**Sol.** Option (D) is correct because Juxtaglomerular cells of kidney secrete erythropoietin hormone which stimulates RBC formation.

Alpha cells of pancreas produce hormone glucagon.

The cells of rostral adenohypophysis synthesizes hormones of anterior lobe of pituitary.

The cells of bone marrow are responsible for formation of formed elements.

## 170. Venereal diseases can spread through :

- Using sterile needles
- Transfusion of blood from infected person
- Infected mother to foetus
- Kissing
- Inheritance

Choose the correct answer from the option given below

- |                           |                           |
|---------------------------|---------------------------|
| (A) (a), (b) and (c) only | (B) (b), (c) and (d) only |
| (C) (b) and (c) only      | (D) (a) and (c) only      |

**Ans. (C)**

**Sol.** Venereal diseases or sexually transmitted diseases or infections are transmitted by sharing of infected needles, surgical instruments with infected person, transfusion of blood or from an infected mother to foetus.

Venereal diseases are not transmitted through kissing or inheritance.

171. Which of the following characteristics is incorrect with respect to cockroach?
- (A) A ring of gastric caeca is present at the junction of midgut and hind gut  
 (B) Hypopharynx lies within the cavity enclosed by the mouth parts  
 (C) In females, 7<sup>th</sup>-9<sup>th</sup> sterna together form a genital pouch  
 (D) 10<sup>th</sup> abdominal segment in both sexes, bears a pair of anal cerci

Ans. (A)

Sol. Option (A) is incorrect because a ring of gastric caecae is present at the junction of foregut and midgut. At the junction of midgut and hindgut, malpighian tubules are present.

Hypopharynx lies within the cavity enclosed by mouthparts.

In female cockroach, the 7<sup>th</sup> sternum is boatshaped and together with the 8<sup>th</sup> and 9<sup>th</sup> sterna forms a genital pouch.

10<sup>th</sup> abdominal segment in both sexes, bears a pair of anal cerci and 9<sup>th</sup> sternum only in male cockroach, bears a pair of chitinous anal style.

172. Match the following:

List-I		List-II	
(a)	<i>Physalia</i>	(i)	Pearl oyster
(b)	<i>Limulus</i>	(ii)	Portuguese Man of War
(c)	<i>Ancylostoma</i>	(iii)	Living fossil
(d)	<i>Pinctada</i>	(iv)	Hookworm

Choose the correct answer from the options given below.

- |          |       |       |      |
|----------|-------|-------|------|
| (a)      | (b)   | (c)   | (d)  |
| (A) (ii) | (iii) | (i)   | (iv) |
| (B) (iv) | (i)   | (iii) | (ii) |
| (C) (ii) | (iii) | (iv)  | (i)  |
| (D) (i)  | (iv)  | (iii) | (ii) |

Ans. (C)

Sol. Option (C) is correct because *Physalia* is commonly known as Portuguese man of war.

*Limulus* is considered as a living fossil and commonly known as king crab.

*Ancylostoma* is a roundworm and commonly known as hookworm.

*Pinctada* is commonly known as pearl oyster, included in phylum Mollusca.

173. Which one of the following organisms bears hollow and pneumatic long bones?
- (A) Neophron (B) Hemidactylus  
 (C) Macropus (D) Ornithorhynchus

Ans. (A)

Sol. Hollow and pneumatic long bones are present in animals that belong to class Aves e.g., Neophron (vulture).

Ornithorhynchus (Platypus) and Macropus (Kangaroo) belong to class Mammalia.

Hemidactylus (Wall lizard) is a member of class Reptilia.

174. The centriole undergoes duplication during:  
(A) S-phase (B) Prophase (C) Metaphase (D) G<sub>2</sub> phase

**Ans.** (A)

**Sol.** During S phase of cell cycle replication of DNA takes place. In animal cells during S phase, centriole duplicates in the cytoplasm.

In G<sub>2</sub> phase there is duplication of mitochondria, chloroplast and Golgi bodies. Tubulin protein is also synthesized during this phase.

During prophase, condensation of chromatin starts.

During metaphase, chromosomes get aligned at equator to form metaphasic plate.

175. During the process of gene amplification using PCR, if very high temperature is not maintained in the beginning, then which of the following steps of PCR will be affected first?

(A) Annealing (B) Extension (C) Denaturation (D) Ligation

**Ans.** (C)

**Sol.** Option (C) is correct. High temperature about 94°C is required for the process of denaturation which is the first step of PCR.

Ligation of DNA fragments is performed with the help of an enzyme called DNA ligase.

Annealing is performed at 50°-60°C which is the second step that can get affected.

Addition of nucleotides to the primer, synthesizing a new DNA strand using only the template sequences with the help of enzyme DNA polymerase is called primer extension/ polymerisation.

176. Which of the following is not an objective of Biofortification in crops?

(A) Improve protein content  
(B) Improve resistance to diseases  
(C) Improve vitamin content  
(D) Improve micronutrient and mineral content

**Ans.** (B)

**Sol.** Biofortification improves vitamin content, protein content and micronutrient and mineral content. It does not create resistance in plants against diseases.

177. Dobson units are used to measure thickness of:

(A) CFCs (B) Stratosphere (C) Ozone (D) Troposphere

**Ans.** (C)

**Sol.** The thickness of the ozone in a column of air from the ground to the top of atmosphere is measured in term of Dobson unit (1 DU = 1ppb).

The lowermost layer of atmosphere is called troposphere.

CFCs are ozone depleting substances. Ozone found in upper part of atmosphere (the stratosphere) is called good ozone.

178. Sphincter of oddi is present at:

(A) Ileo-caecal junction  
(B) Junction of hepato-pancreatic duct and duodenum  
(C) Gastro-oesophageal junction  
(D) Junction of jejunum and duodenum

**Ans. (B)**

**Sol.** The bile duct and the pancreatic duct open together into the duodenum as the common hepato-pancreatic duct which is guarded by a sphincter called the sphincter of Oddi.

Ileo-caecal valve is present at the junction of ileum and caecum to prevent the backflow of faecal matter into the ileum in humans.

Gastro-oesophageal sphincter regulates the opening of oesophagus into stomach.

**179.** Select the favourable conditions required for the formation of oxyhaemoglobin at the alveoli.

- (A) High  $pO_2$ , low  $pCO_2$ , less  $H^+$ , lower temperature
- (B) Low  $pO_2$ , high  $pCO_2$ , more  $H^+$ , higher temperature
- (C) High  $pO_2$ , high  $pCO_2$ , less  $H^+$ , higher temperature
- (D) Low  $pO_2$ , low  $pCO_2$ , more  $H^+$ , higher temperature

**Ans. (A)**

**Sol.** The factors favourable for the formation of oxyhaemoglobin at the alveolar level are; high  $pO_2$ , low  $pCO_2$ , less  $H^+$  concentration and lower temperature.

The conditions favourable for the dissociation of oxygen from oxyhaemoglobin at the tissue level are; low  $pO_2$ , high  $pCO_2$ , high  $H^+$  concentration and high temperature.

**180.** Identify the incorrect pair

- |               |   |                |
|---------------|---|----------------|
| (A) Alkaloids | - | Codeine        |
| (B) Toxin     | - | Abrin          |
| (C) Lectins   | - | Concanavalin A |
| (D) Drugs     | - | Ricin          |

**Ans. (D)**

**Sol.** Option (D) is incorrect because ricin is a toxin obtained from Ricinus plant. Vinblastin and curcumin are drugs.

Morphine and codeine are alkaloids.

Abrin is also a toxin obtained by plant Abrus.

Concanavalin A is a lectin.

**181.** Which of the following statements wrongly represents the nature of smooth muscle?

- (A) These muscle have no striations
- (B) They are involuntary muscles
- (C) Communication among the cells is performed by intercalated discs
- (D) These muscles are present in the wall of blood vessels

**Ans. (C)**

**Sol.** Option (C) is incorrect because intercalated discs are found only in cardiac muscle tissue.

Smooth muscle fibres are non-striated and involuntary in nature and are present in the wall of blood vessels, uterus, gall bladder, alimentary canal etc.

**182.** For effective treatment of the disease, early diagnosis and understanding its pathophysiology is very important. Which of the following molecular diagnostic techniques is very useful for early detection?

- (A) Western Blotting Technique
- (B) Southern Blotting Technique

- (C) ELISA Technique  
(D) Hybridization Technique

**Ans. (B/C\*)**

**Sol.** ELISA can be used for early detection of an infection either by detecting the presence of pathogenic antigen or by detecting the antibodies synthesized against the pathogen.

Option (B) Southern blotting is used to detect a specific DNA sequence in the given sample and can be detected prior to antibody formation. One can detect presence of pathogenic DNA/RNA.

In hybridization technique a ssDNA/ssRNA tagged with a radioactive molecule (probe) is allowed to hybridize its complementary DNA in a clone of cells followed by detection using autoradiography. It is used to find a mutated gene.

Western blotting technique is used to detect a specific protein molecule among a mixture of proteins.

**183.** Match List-I with List-II.

List-I		List-II	
(a)	Vaults	(i)	Entry of sperm through Cervix is blocked
(b)	IUDs	(ii)	Removal of Vas deferens
(c)	Vasectomy	(iii)	Phagocytosis of sperms within the Uterus
(d)	Tubectomy	(iv)	Removal of fallopian tube

Choose the correct answer from the option given below

- |           |       |       |       |
|-----------|-------|-------|-------|
| (a)       | (b)   | (c)   | (d)   |
| (A) (iv)  | (ii)  | (i)   | (iii) |
| (B) (i)   | (iii) | (ii)  | (iv)  |
| (C) (ii)  | (iv)  | (iii) | (i)   |
| (D) (iii) | (i)   | (iv)  | (ii)  |

**Ans. (B)**

**Sol.** Diaphragms, cervical caps and vaults are barrier methods of contraception for female which works by blocking the entry of sperms through the cervix.

IUDs increase phagocytosis of sperms within the uterus.

Vasectomy is a surgical method of contraception in males in which a small part of the vas deferens is removed or tied up through a small incision on the scrotum.

Tubectomy is a surgical method of contraception in females where a small part of the fallopian tube is removed or tied up through a small incision in the abdomen or through vagina.

**184.** The organelles that are included in the endomembrane system are

- (A) Endoplasmic reticulum, Mitochondria, Ribosomes and Lysosomes  
(B) Endoplasmic reticulum, Golgi complex, Lysosomes and Vacuoles  
(C) Golgi complex, Mitochondria, Ribosomes and Lysosomes  
(D) Golgi complex, Endoplasmic reticulum, Mitochondria and Lysosomes

**Ans. (B)**

**Sol.** Endomembrane system consist of endoplasmic reticulum, Golgi complex, vacuoles and lysosomes.

Mitochondria is semi-autonomous cell organelle.

Ribosome is non-membranous cell organelle.

185. Which stage of meiotic prophase shows terminalisation of chiasmata as its distinctive feature?  
(A) Leptotene (B) Zygotene (C) Diakinesis (D) Pachytene

Ans. (C)

Sol. In meiosis I, chiasmata (X shaped structure) is formed in diplotene stage while it terminalise in diakinesis stage.

Bivalents are formed in zygotene stage and crossing over takes place in pachytene stage.

Compaction of chromosomal material occurs in leptotene stage.

### SECTION - B

186. Which of these is not an important component of initiation of parturition in humans ?  
(A) Increase in estrogen and progesterone ratio  
(B) Synthesis of prostaglandins  
(C) Release of Oxytocin  
(D) Release of Prolactin

Ans. (D)

Sol. At the end of gestation, the completely developed foetus is expelled out. This process is called parturition.

Parturition is controlled by a complex neuroendocrine mechanism.

Estrogen and progesterone ratio increases as estrogen levels rise significantly.

Prostaglandins, which stimulate uterine contractions are also produced that act on myometrium.

Oxytocin, the main hormone, also called as birth hormone is released by maternal pituitary, which brings about strong uterine contractions.

Prolactin is a lactation hormone that has no role in initiation of parturition.

187. Which of the following is not a step in Multiple Ovulation Embryo Transfer Technology (MOET)?  
(A) Cow is administered hormone having LH like activity for super ovulation  
(B) Cow yields about 6-8 eggs at a time  
(C) Cow is fertilized by artificial insemination  
(D) Fertilized eggs are transferred to surrogate mothers at 8-32 cell stage

Ans. (A)

Sol. Multiple Ovulation Embryo Transfer Technology is used for herd improvement in short time.

Cows are administered hormones, with FSH-like activity for superovulation.

8-32 celled embryos are transferred to surrogate mothers.

6-8 eggs are produced per cycle.

Cows can be fertilised by artificial insemination.

188. Match List-I with List - II

List - I		List - II	
(a)	Allen's Rule	(i)	Kangaroo rat
(b)	Physiological adaptation	(ii)	Desert lizard
(c)	Behavioural adaptation	(iii)	Marine fish at depth
(d)	Biochemical adaptation	(iv)	Polar seal

Choose the correct answer from the options given below.

- |          |       |       |       |
|----------|-------|-------|-------|
| (a)      | (b)   | (c)   | (d)   |
| (A) (iv) | (ii)  | (iii) | (i)   |
| (B) (iv) | (i)   | (ii)  | (ii)  |
| (C) (iv) | (i)   | (ii)  | (iii) |
| (D) (iv) | (iii) | (ii)  | (i)   |

**Ans.** (C)

**Sol.** Polar seal generally has shorter ears and limbs (extremities) to minimise heat loss. This is with reference to Allen's rule.

Kangaroo rat exhibits physiological adaptation.

Desert lizard shows behavioural adaptation.

They lack the physiological ability to cope-up with extreme temperature but manage the body temperature by behavioural means.

Marine fishes at depth are adapted biochemically to survive in great depths in ocean.

**189.** Assertion (a): A person goes to high altitude and experiences 'altitude sickness' with symptoms like breathing difficulty and heart palpitations.

Reason (r): Due to low atmospheric pressure at high altitude, the body does not get sufficient oxygen.

In the light of the above statements, choose the correct answer from the options given below

- (A) Both (a) and (r) are true and (r) is the correct explanation of (a)  
 (B) Both (a) and (r) are true but (r) is not the correct explanation of (a)  
 (C) (a) is true but (r) is false  
 (D) (a) is false but (r) is true

**Ans.** (A)

**Sol.** Altitude sickness can be experienced at high altitude where body does not get enough oxygen due to low atmospheric pressure and causes nausea, fatigue and heart palpitations.

Hence correct option is (A) as [r] is correct explanation of [a].

**190.** Following are the statements with reference to 'lipids'.

- (a) Lipids having only single bonds are called unsaturated fatty acids  
 (b) Lecithin is a phospholipid.  
 (c) Trihydroxy propane is glycerol.  
 (d) Palmitic acid has 20 carbon atoms including carboxyl carbon.  
 (e) Arachidonic acid has 16 carbon atoms. Choose the correct answer from the options given below.

- (A) (a) and (b) only                      (B) (c) and (d) only  
 (C) (b) and (c) only                      (D) (b) and (e) only

**Ans. (C)**

**Sol.** The correct option is (C) because lipids having only single bonds are called saturated fatty acids and lipids having one or more C = C double bonds are called unsaturated fatty acids.

Palmitic acid has 16 carbon atoms including carboxyl carbon.

Arachidonic acid has 20 carbon atoms including the carboxyl carbon.

Lecithin is a phospholipid found in cell membrane.

Glycerol has 3 carbons, each bearing a hydroxyl (-OH) group.

**191.** Match List-I with List-II

List-I		List-II	
(a)	Scapula	(i)	Cartilaginous joints
(b)	Cranium	(ii)	Flat bone
(c)	Sternum	(iii)	Fibrous joints
(d)	Vertebral column	(iv)	Triangular flat bone

Choose the correct answer from the options given below

- |          |       |       |      |
|----------|-------|-------|------|
| (a)      | (b)   | (c)   | (d)  |
| (A) (i)  | (iii) | (ii)  | (iv) |
| (B) (ii) | (iii) | (iv)  | (i)  |
| (C) (iv) | (ii)  | (iii) | (i)  |
| (D) (iv) | (iii) | (ii)  | (i)  |

**Ans. (D)**

**Sol.** The correct option is (D).

Scapula is a large triangular flat bone situated in the dorsal part of the thorax between the second and the seventh ribs.

Fibrous joint is shown by the flat skull bones which fuse end-to-end with the help of dense fibrous connective tissues in the form of sutures, to form the cranium.

Sternum is a flat bone on the ventral midline of thorax.

Cartilaginous joints between the adjacent vertebrae in the vertebral column permits limited movements.

**192.** Identify the types of cell junctions that help to stop the leakage of the substances across a tissue and facilitation of communication with neighbouring cells via rapid transfer of ions and molecules.

- (A) Gap junctions and Adhering junctions, respectively  
 (B) Tight junctions and Gap junctions, respectively  
 (C) Adhering junctions and Tight junctions, respectively.  
 (D) Adhering junctions and Gap junctions, respectively

**Ans. (B)**

**Sol.** Three types of junctions are found in tissues

Tight junctions stop leakage of substances from leaking across a tissue.

Adhering junctions cement and keep neighbouring cells together.

Gap junctions or communication junctions facilitate communication between cells by connecting the cytoplasm of adjoining cells.

- 193.** Statement I: The codon 'AUG' codes for methionine and phenylalanine.  
Statement II: 'AAA' and 'AAG' both codons code for the amino acid lysine.  
In the light of the above statements, choose the correct answer from the options given below.
- (A) Both Statement I and Statement II are true  
(B) Both Statement I and Statement II are false  
(C) Statement I is correct but Statement II is false  
(D) Statement I is incorrect but Statement II is true

**Ans. (D)**

**Sol.** AUG has dual functions, it codes for methionine. It also acts as initiator codon. AUG does not code for phenylalanine. Statement II is true.

- 194.** Which of the following secretes the hormone, relaxin, during the later phase of pregnancy?  
(A) Graafian follicle (B) Corpus luteum (C) Foetus (D) Uterus

**Ans. (B)**

**Sol.** The hormone relaxin is produced in the later phase of pregnancy. It is produced by the ovary. Graafian follicle is not formed when the woman is pregnant. Uterus and foetus do not produce relaxin. Relaxin is produced by the corpus luteum present in the ovary. Ruptured Graafian follicle is called corpus luteum, which has endocrine function.

- 195.** Following are the statements about prostomium of earthworm.
- (a) It serves as a covering for mouth.  
(b) It helps to open cracks in the soil into which it can crawl.  
(c) It is one of the sensory structures.  
(d) It is the first body segment.

Choose the correct answer from the options given below.

- (A) (a), (b) and (c) are correct (B) (a), (b) and (d) are correct  
(C) (a), (b), (c) and (d) are correct (D) (b) and (c) are correct

**Ans. (A)**

**Sol.** The anterior end of the earthworm has mouth which has covering called prostomium. Prostomium acts as a wedge to force open cracks in the soil. Prostomium has receptors, so it is sensory in function. The first body segment of earthworm is the peristomium

- 196.** Which one of the following statements about Histones is wrong?  
(A) Histones are organized to form a unit of 8 molecules  
(B) The pH of histones is slightly acidic  
(C) Histones are rich in amino acids - Lysine and Arginine  
(D) Histones carry positive charge in the side chain

**Ans. (B)**

**Sol.** Histones are rich in basic amino acids residue lysine and arginine with charged side chain.

There are five types of histone proteins i.e.,  $H_1, H_2A, H_2B, H_3$  and  $H_4$ . Four of them occur in pairs to produce a unit of 8 molecules (histone octamer) The pH of histones is basic.

197. During muscular contraction which of the following events occur?

- (a) 'H' zone disappears
- (b) 'A' band widens
- (c) 'I' band reduces in width
- (d) Myosine hydrolyzes ATP, releasing the ADP and Pi.
- (e) Z-lines attached to actins are pulled inwards.

Choose the correct answer from the options given below:

- (A) (a), (c), (d), (e) only
- (B) (a), (b), (c), (d) only
- (C) (b), (c), (d), (e) only
- (D) (b), (d), (e), (a) only

Ans. (A)

Sol. The correct option is (A) because the length of A-band is retained. During muscle contraction, the following events occur:

- (1) The globular head of myosin acts as ATPase and hydrolyses ATP molecule and eventually leads to the formation of cross bridge.
- (2) This pulls the actin filament towards the centre of 'A-band'.
- (3) The Z-line attached to these actins are also pulled inwards thereby causing a shortening of the sarcomere.
- (4) The thin myofilaments move past the thick myofilaments due to which the H-zone narrows. This reduces the length of I-band but retains the length of A-band.
- (5) The myosin then releases ADP + Pi, and goes back to its relaxed state.

198. The Adenosine deaminase deficiency results into

- (A) Dysfunction of Immune system
- (B) Parkinson's disease
- (C) Digestive disorder
- (D) Addison's disease

Ans. (A)

Sol. Adenosine deaminase (ADA) enzyme is crucial for the immune system to function. Hence, its deficiency results in the dysfunction of immune system.

Hyposecretion of hormones of the adrenal cortex causes Addison's disease.

Parkinson's disease is a long-term degenerative disorder of the central nervous system.

Disorders which affect GIT & associated glands are called digestive disorders.

199. Match List - I with List - II

List - I		List - II	
(a)	Adaptive radiation	(i)	Selection of resistant varieties due to excessive use of herbicides and pesticides
(b)	Convergent evolution	(ii)	Bones of forelimbs in Man and Whale
(c)	Divergent evolution	(iii)	Wings of Butterfly and Bird
(d)	Evolution by anthropogenic action	(iv)	Darwin Finches

Choose the correct answer from the options given below.

(a)	(b)	(c)	(d)
(A) (iv)	(iii)	(ii)	(i)
(B) (iii)	(ii)	(i)	(iv)
(C) (ii)	(i)	(iv)	(iii)
(D) (i)	(iv)	(iii)	(ii)

**Ans. (A)**

**Sol.** The correct option is (A)

Adaptive radiation is the process of evolution of different species in a given geographical area starting from a point and literally radiating to other areas of geography, for example : Darwin's finches.

Analogous organs which are not anatomically similar structures though they perform similar functions, are a result of convergent evolution, for example: Wings of butterfly and of birds.

Homologous organs which are anatomically similar structures but perform different functions according to their needs, are a result of divergent evolution, for example : Bones of forelimbs in man and whale.

Evolution by anthropogenic action means evolution due to human interference, for example: Antibiotic resistant microbes, herbicides resistant varieties and pesticide resistant varieties.

200. Match List-I with List-II

List -I		List -II	
(a)	Filariasis	(i)	<i>Haemophilus influenzae</i>
(b)	Amoebiasis	(ii)	<i>Trichophyton</i>
(c)	Pneumonia	(iii)	<i>Wuchereria bancrofti</i>
(d)	Ringworm	(iv)	<i>Entamoeba histolytica</i>

Choose the correct answer from the options given below

(a)	(b)	(c)	(d)
(A) (iv)	(i)	(iii)	(ii)
(B) (iii)	(iv)	(i)	(ii)
(C) (i)	(ii)	(iv)	(iii)
(D) (ii)	(iii)	(i)	(iv)

**Ans. (B)**

**Sol.** The correct option is (B).

Filariasis is the disease caused by *Wuchereria bancrofti*, filarial worm.

Amoebiasis/Amoebic dysentery is caused by a protozoan parasite *Entamoeba histolytica* in the large intestine of human.

Pneumonia is caused by bacteria like *Streptococcus pneumoniae* and *Haemophilus influenzae*.

Ringworm is caused by fungi belonging to genera *Microsporum*, *Trichophyton* and *Epidermophyton*.

